

## **Growth of marine ectotherms is regionally constrained and asymmetric with latitude**

Reed, Adam J.; Godbold, Jasmin; Grange, Laura; Solan, Martin

### **Global Ecology and Biogeography**

DOI:

<https://doi.org/10.1111/geb.13245>

Published: 01/03/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Reed, A. J., Godbold, J., Grange, L., & Solan, M. (2021). Growth of marine ectotherms is regionally constrained and asymmetric with latitude. *Global Ecology and Biogeography*, 30(3), 578-589. <https://doi.org/10.1111/geb.13245>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Growth of marine ectotherms is regionally constrained and  
asymmetric with latitude**

Journal:	<i>Global Ecology and Biogeography</i>
Manuscript ID	GEB-2019-0513.R4
Manuscript Type:	Meta-Analysis
Keywords:	biogeography, climate change, growth, macrophysiology, regional constraints, resource management

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Title:** Growth of marine ectotherms is regionally constrained and asymmetric with latitude

**Running Title:** Global growth of marine ectotherms

For Peer Review

**Abstract****Aim**

Growth rates of organisms are routinely used to summarise physiological performance, but the consequences of local evolutionary history and ecology are largely missed by analyses on wide biogeographic scales. This broad approach has been commonly applied to other physiological parameters across terrestrial and aquatic environments. Here, we examine growth rates of marine bivalves across all biogeographic realms, latitude, and temperature, with analyses to determine regional effects on growth on global scales.

**Location**

Global: Marine Ecosystems

**Time Period**

1930–2018

**Major Taxa**

Bivalves

**Methods**

We use a comprehensive data-set of bivalve growth parameters ( $n=966$ , 243 species) representing all biogeographic realms to calculate overall growth performances. We use these data with environmental temperature to analyse global patterns in growth, accounting for regional primary productivity and phylogeny using general additive mixed and linear models. The Arrhenius relationship and corresponding activation energies are used to quantify the sensitivity to temperature in each biogeographic realm and province.

**Results**

Our analyses show that bivalve growth demonstrates latitudinal asymmetry and exhibits non-linear relationships with latitude. We find that overall growth performance is affected by temperature and particulate organic carbon, but the form of these relationships differ with phylogeny. Growth is slower in the Antarctic and more sensitive to increasing temperature than in the Arctic, and decrease with increasing temperature in some tropical realms, a previously unidentified and fundamental difference in growth and physiological sensitivity.

**Main Conclusions**

Our findings provide compelling evidence that the widely used curvilinear relationship between temperature and growth rates in marine ectotherms is an inappropriate descriptor of thermal sensitivity, because it normalises regional variations in physiological performance. Without a more detailed assessment of global physiological patterns, the responses of species to local variations associated with climate change will be under-appreciated in global assessments of climate risk, minimising the effectiveness of management and conservation.

**Keywords:** biogeography, climate change, growth, macroecology, physiology, regionally constrained, resource management

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Introduction**

Air and sea temperatures are rising globally, and modelled projections indicate that this trend will continue leading to significant risks of extinction in terrestrial and aquatic systems (Thomas et al., 2004; Urban, 2015), that are likely to disproportionately affect ectothermic (Deutsch et al., 2008; Ohlberger, 2013) and marine organisms (Pinsky, Eikeset, McCauley, Payne, & Sunday, 2019). However, global warming is not uniformly expressed (Brierley & Kingsford, 2009; Seneviratne et al., 2018), and observed ecological responses often reflect the thermal range and physiology of species (Burrows et al., 2011; Hoegh-Guldberg & Bruno, 2010; Root et al., 2003) rather than the expectation derived from the pooling of laboratory based studies that focus on acute thermal shock response (Peck, Webb, & Bailey, 2004; Pörtner, 2001). Further, some fundamental physiological assumptions, such as variation in lifespan (Moss et al., 2016), metabolic rates (Heilmayer, Brey, & Pörtner, 2004), growth (Pörtner, Storch, & Heilmayer, 2005), and acclimation capacity (Seebacher, White, & Franklin, 2014) over wide geographical ranges are based on data with poor spatial resolution and/or are fitted with curvilinear models that do not adequately account for variation in regional environmental conditions. However, incorporation of local processes is vital, as environmental history and setting affects how individuals respond to altered environmental conditions (Calosi, De Wit, Thor, & Dupont, 2016; Gladstone-Gallagher, Pilditch, Stephenson, & Thrush, 2019; Deutsch, Penn & Seibel, 2020) and affect ecosystem functioning (Godbold & Solan, 2013; Wohlgemuth, Solan, & Godbold, 2016).

The Arrhenius relationship defines how increasing temperature accelerates metabolic processes by speeding up reaction rates (Clarke, 2017), and derived activation energies quantify the sensitivity of the response measured (Peck, 2018). This relationship of log rate against inverse temperature produces a straight line, and is often used to fit the latitudinal temperature gradient to facilitate understanding of whole animal physiology (Heilmayer, Brey & Pörtner, 2004; Peck, 2018), yet departures from the projected relationship have been identified (Peck, 2016) and have typically been explained by constraints imposed by subtle ecological distinctions (Deutsch et al., 2008), evolutionary history (Moss, Ivany, Silver, Schue, & Artruc, 2017), or molecular constraints (Peck, 2016). However, these deviations are integral to the determination of relevant ecosystem responses to climate change (Godbold & Solan, 2013), and are especially important in regions where human well-being is highly dependent on ecosystem services (van der Schatte Olivier et al., 2018).

Measuring physiological variation across large spatial scales provides a definitive means to understand species-specific physiological responses to a dynamic environment, but is logistically challenging and not commonly achieved in aquatic systems (Osovitz & Hofmann, 2007). A pragmatic alternative is to use growth rates in natural populations (Reed, Linse, & Thatje, 2014) and, as growth is a trade-off with metabolic rates and reproduction, it represents a reasonable approximation of whole animal physiology (Clarke, 2003; Pörtner et al., 2001). Here, we use overall growth performance (OGP, the point of inflection in a von Bertalanffy growth curve; Brey, 1999) in marine bivalves to quantify growth constraints between biogeographical realms. Marine bivalves are globally distributed and their growth is integrated within their shells (Moss et al., 2016). OGP is preferential to the more frequently used growth constant ( $k$ ) as it takes into consideration the theoretical maximum

shell length of individuals, allowing comparisons between taxa (Brey, 1999). By calculating activation energies in biogeographical realms we quantify the thermal sensitivity of regional growth and identify realms at greater risk to projected temperature change based on how local species growth differs across local temperature ranges. We anticipate that regional disparities will expose a fundamental misunderstanding of physiological responses in regions at the upper and lower thermal limits of these species, especially where previous efforts have pooled data or not taken into account evolutionary history or unique local ecologies. By exploring the spatiotemporal distribution of physiological characteristics across the globe (Osovit & Hofmann, 2007), physiological projections within distinct regions can be determined (Chown, Sinclair, Leinaas, & Gaston, 2004; Pörtner & Knust, 2007; Somero, 2010), shifting paradigms that are likely to benefit conservation and management efforts (Stuart-Smith, Edgar, Barrett, Kininmonth, & Bates, 2015).

## Methods

We searched the Thomson Reuters Web of Science collection (<http://www.webofknowledge.com>, accessed July, 2019) using the search terms “(“bivalve”, “growth”, and “von Bertalanffy”)” in the titles, keywords and abstracts of all document types from 1950-2018. Further relevant publications were obtained by manually checking references cited by the authors of the returns from our search. We excluded all publications that were based on data from cultured or artificially reared populations, but data from PhD theses or ‘grey literature’ were included after critical examination of the methodology to ensure consistency with published growth parameters. A list of the data sources is found in Appendix S1.

### Growth data

Citation returns were manually searched for reported values of parameters of growth from von Bertalanffy growth functions ( $k$ ,  $L_{\infty}$ , and, if possible,  $L_T$ ). A total of 429 peer-reviewed publications revealed 966 growth parameters for 243 species of bivalve from 143 Genera, 44 families and three subclasses (Reed et al. 2020). As well as taxonomic information, geographical location, depth, and temperature were extracted when available (see *Environmental Data*). When the location of a study was not provided, latitude and longitude coordinates and/or water depth were retrieved from Google Earth (<http://earth.google.com/>) and manually cross referenced with site descriptions within the source publication. Taxonomy was verified and updated to latest classification using the World Online Register of Marine Species (WoRMS Editorial Board, 2020)

Where specific size-at-age values were not presented in the original publication and had to be derived, values were extracted from graphical summaries using Web Plot Digitiser (<https://automeris.io/WebPlotDigitizer/>). We determined the required growth data using these data by applying the von Bertalanffy model as described by Brey 1999:

$$L_t = L_{\infty}(1 - e^{-k(t-t_0)})$$

where  $k$  is the growth constant and  $L_{\infty}$  is the asymptotic length.

1  
2  
3 153 In contrast to previous publications on global growth data, we were able to calculate Overall Growth  
4 154 Performance (OGP) as we only included publications with growth parameters from a von Bertalanffy  
5 155 growth curve. Previous global studies comparing growth have only used single parameters directly  
6 156 derived from the growth curve and typically use  $k$  (The Brody growth coefficient) as a measure of  
7 157 growth between species (Moss et al., 2016; Peck, 2016). However, growth is non-linear and it is not  
8 158 appropriate to compare growth using single parameters from the growth function in a statistical  
9 159 analysis (Brey, 1999). OGP derived from the Bertalanffy growth function makes growth comparable  
10 160 between populations and species by removing individual variation and is defined as the point of  
11 161 inflection on the Bertalanffy growth curve (Brey, 1999). Overall Growth Performance is calculated  
12 162 from the equation:  
13 163

14 164 
$$OGP = \log(k L_{\infty})$$

15 165 *Environmental data*

16 166 When temperature data was not reported, bottom water temperatures were extracted from Ocean  
17 167 Data View v. 5.1.0 and the World Ocean Atlas 2013 database (Locarnini et al., 2013) at a resolution  
18 168 of 0.25 degrees and using coordinates from the corresponding paper. We only used recorded bottom  
19 169 temperatures and not sea surface temperatures which are commonly used in global marine studies,  
20 170 as the bottom temperatures are more representative of the benthic environment.

21 171 OGP data was grouped based on Spalding's classification of biogeographical latitude zones, realms,  
22 172 and provinces (Spalding et al., 2007). While all biogeographic realms are represented in our data-set,  
23 173 provinces with no samples are not included in our maps and are shown as data deficient gaps. Unlike  
24 174 previous studies examining physiological patterns across wide geographic areas, we differentiate  
25 175 OGP between the different biogeographical latitude zones, realms, and provinces which allows a  
26 176 higher resolution and critical examination of regional OGP.

27 177 POC flux to the seafloor was derived by applying a vertical flux attenuation equation to satellite-  
28 178 derived export fluxes. The export flux is calculated from satellite-derived primary production (Carr,  
29 179 2001) and sea surface temperature using the relationship defined in Henson et al. (2011). Flux  
30 180 attenuation is described by "Martin's b" parameter (Martin et al., 1987) and is estimated globally using  
31 181 a collation of deep moored sediment trap data as described in Henson et al. (2012). Seafloor depth is  
32 182 taken from the ETOP01 global relief model (Amante and Eakins, 2009). All data are spatially  
33 183 averaged onto a 1x1 degree grid prior to analysis. POC flux to the seafloor (in  $\text{gC m}^{-2} \text{yr}^{-1}$ ) is extracted  
34 184 from the global data at each sample location using a nearest neighbour approach. The POC flux to  
35 185 seafloor could not be estimated at several shallow shelf locations due to the limitations of satellite-  
36 186 derived data and the flux model in very shallow waters. In these instances, the nearest geographical  
37 187 point (linear Euclidean distance) with data was supplemented.

38 188 *Arrhenius Relationship*

39 189 The Arrhenius relationship relates chemical reaction rates to temperature, and is expressed by;



$$k = Ae^{\left(\frac{-E_a}{RT}\right)}$$

Where  $k$  is the rate constant,  $R$  is the gas constant,  $T$  is absolute temperature, and  $E_a$  is the activation energy. Physiological processes have been proven to respond predictably by following the Arrhenius relationship within normal biological temperature ranges (Clarke & Johnston, 2003; Heilmayer et al., 2004). We calculate activation energies (ev) from the linear relationship between absolute temperature and overall growth performance to represent the sensitivity of OGP to temperature. These values can be used to directly compare biological reaction rates and processes such as growth. For biological processes activation energies can range between 0.2 and 1.2ev (Gillooly, Brown, West, Savage, & Charnov, 2001) but usually vary between 0.6 and 0.7ev (Dell, Pawar, & Savage, 2011).

### *Statistical data analysis*

To determine the direction and significance of the relationship between OGP and latitudinal zone between Realms, we used a generalised additive model (GAM) with a Gaussian distribution using “REML” estimation. We further account for regional differences in temperature and primary productivity (POC  $\text{gC m}^{-2} \text{yr}^{-1}$ ;  $\ln$  transformed) and test for any differences in growth rates of related species occurring at similar temperatures but in different regions (Subclass, 2 levels: Heterodonta and Pterimorphia; the subclass Protobranchia (9 observations) were excluded from the analysis). To investigate the relationship between OGP, temperature and  $\ln(\text{POC})$  we used a Generalized Additive Mixed Model (GAMM) incorporating the factor “Realm” as a random effect. To estimate the optimal amount of smoothing in both analyses ( $s(\text{latitude})$  and  $s(\text{temperature})$ ,  $s(\ln(\text{POC}))$ ) we used cross-validation (Zuur, Ieno, Walker, Saveliev, & Smith, 2009; Zuur, Saveliev, Ieno, 2015) followed by optimisation of  $k$  following Wood et al. (2017). Smoothers for Latitude and  $\ln(\text{POC})$  were fitted with cubic spline regression ( $cs$ ), whilst the smoother for temperature was fitted with thin plate regression ( $tp$ ). For both analyses final best models were based on Akaike information criterion (AIC) and residual fits were examined (Zuur, Saveliev, Ieno, 2015; Wood et al. 2017). Predicted values and 95% Confidence Intervals of final models were determined and fitted following Zuur, Saveliev & Ieno (2015).

To determine the importance of latitudinal zone and the Arrhenius model on  $\ln(\text{OGP})$  we developed an ANCOVA model containing latitudinal zone as a nominal explanatory variable with 4 levels (Antarctic, Arctic, Temperate and Tropical) and the Arrhenius model represented by inverse temperature (Kelvins) as a continuous explanatory variable. We visually assessed model assumptions (homogeneity of variances and normality) which revealed patterns in the residual spread. To model the heteroscedasticity in the variance structure we incorporated the variable “latitude” as a variance covariate using  $\text{varIdent}$  (Pinheiro & Bates, 2000).

Latitude and longitude coordinates of the individual data points were used to visually assess the effects of spatial autocorrelation using bubble plots (Pebesma & Graeler 2019) and variograms (Zuur et al., 2009). Analyses were conducted in R (R Development Core Team 2018) using the “nlme”



library for the extended linear model (Pinheiro, Bates, DebRoy, & Sarkar, 2018), the “mgcv” library for the additive (mixed) models (Wood, 2019) and “gstat” for investigating spatial autocorrelation (Pebesma et al., 2019).

## Results

We established 966 measurements of bivalve growth parameters spanning 243 species and representing all 12 biogeographic realms. Measurements of overall growth performance (OGP) ranged from 0.01 to 2.68 and extended from 80 °N to 77 °S and from 176 °W to 175 °E. Depth was rarely reported in the literature, however most growth measurements were from species known from the coast or shelf (<200 m) regions with 8 species specifically reported from deeper waters (250 – 4600 m) and no observations from hadal depths. The distribution of data corresponds mostly to the Temperate North Atlantic Realm, (n = 367), followed by the Temperate North Pacific (n = 173) and Temperate South America (n = 148) Realms, and indicates that most observations emanate from the northern hemisphere (73.3%). Temperate Latitude Zones are highly represented (n = 737), followed by Tropical (n = 124) and Polar (n = 105) Latitude Zones.

Using these data, our analyses reveal that OGP in marine bivalves increases with decreasing latitude and, for the first time, we show that the form of this relationship is non-linear (GAM, edf= 8.651, F = 13.927 p<0.0001, Figure 1a) and that OGP differs between Realms (GAM, F = 10.13, p <0.0001, Figure 1b). Specifically, we find that there is an increase in mean OGP in the North Atlantic above 50°N driven by species in the Northern European Seas, whilst the Mediterranean Sea is characterised by lower mean OGP ( $1.14 \pm 0.31$ , Figure 1c, Figure S2.1). In the Southern Hemisphere, the highest mean OGP ( $\pm$  95% CI) is observed in the tropical regions of the Central and East Indo-Pacific ( $1.93 \pm 0.16$  and  $1.67 \pm 0.20$ , respectively), decreasing towards the temperate realms and into the Southern Ocean ( $0.57 \pm 0.17$ ). In the Northern Hemisphere, the results indicate a second peak in OGP (Figure 1a) at around 50°N in the Temperate North Pacific and North Atlantic realms. Between latitudinal zones, mean OGP ( $\pm$  95% CI) was lowest in the Antarctic and Arctic ( $0.575 \pm 0.173$  and  $0.930 \pm 0.073$ , respectively) and highest in Tropical areas ( $1.569 \pm 0.078$ ) (Figure 1a), albeit influenced by a high mean OGP in the Central and East Indo-Pacific realms (Figure 1b).

Our analyses confirm a positive relationship between temperature and decreasing latitude (Figure S2.2). Whilst OGP increases with temperature (GAMM, edf=7.821, F = 14.21, p<0.0001), we find a non-linear, stepwise relationship (Figure 2a) that broadly corresponds to Polar (-2 to 6°C), Temperate (7 to 17°C), and Tropical (18 to 27°C) latitudinal zones. Furthermore, the requirement to incorporate Realm as a random effect confirms the important role of biogeographical divisions in determining temperature dependent OGP. We found that particulate organic carbon levels are highest in temperate latitude zones and overlap with the Arctic, whilst are low throughout the Antarctic and tropical areas (Figure S2.3), but that the shape of the relationship between OGP and  $\ln(\text{POC})$  differs between the two Subclasses of bivalves and is only significant for the Pteriomorphia ( $\ln(\text{POC})$  : Pteriomorphia: edf = 5.9, F = 10.142, p<0.0001; Heterodonta: edf = 1, F = 1.506, p = 0.22). For the Pteriomorphia (8 families, n = 408) there is a peak in OGP at intermediate levels of  $\ln(\text{POC})$ , whilst for

the Heterodonta (29 families,  $n = 549$ ) there is no difference in OGP with changing  $\ln(\text{POC})$  (Figure 2b). Overall, OGP is significantly higher in the Pteriomorphia (mean =  $1.409 \pm 0.047$ ,  $n = 407$ ) than the Heterodonta (mean =  $1.228 \pm 0.032$ ,  $n = 550$ ; GAMM,  $t = 11.55$ ,  $p < 0.0001$ ; Figure 2c., Figure S2.4; S2.5).

We fitted an Arrhenius model using inverse temperature ( $1000/T$  in Kelvins) and  $\ln(\text{OGP})$ , and their interaction, to test OGP against the expected physiological relationship with temperature (Figure 3a) We find that  $\ln(\text{OGP})$  is positively affected by the independent effects of latitude (GLS, L. ratio = 33.561, d.f. = 1,  $p < 0.0001$ ) and inverse temperature (GLS, L. ratio = 31.967, d.f. = 3,  $p < 0.0001$ ), but deviate from the expected relationship in the Antarctic at temperatures  $< 0^\circ\text{C}$  (Figure 3b). The activation energies which quantify the Arrhenius relationship within each biogeographical Latitudinal Zone and Realm (Figure 4) reveal high sensitivity to temperature in the Antarctic Latitude Zone (0.987 ev) and low sensitivity across the Tropical Latitude Zone (0.035 ev) (Figure 4a; Figure S2.6). However, between the biogeographical realms the heterogeneity within tropical and polar zones becomes more apparent, with values between -0.770 ev and 0.987 ev (Figure 4b; Figure S2.7). These data show a negative relationship of OGP with increasing temperature in East Indo-Pacific, West Indo-Pacific, Tropical Pacific and Temperate Australia, indicating reduced growth in species in these regions with increasing temperatures, while all temperate and polar biogeographic realms show a positive relationship with temperature.

## Discussion

This global database of overall growth performance (OGP) in marine bivalves confirms that growth increases with decreasing latitude (Moss et al., 2016; Pörtner, Storch, & Heilmayer, 2005), but we reveal that the form of this relationship is non-linear and depends on biogeographical context. Latitudinal variation of physiological parameters has previously been associated with seasonality and genetic adaptations to a specific temperature range (Yamahira & Conover, 2002), although temperature dependent hypoxia may also have a major role in determining biogeographical patterns (Deutch, Penn, & Seibel, 2020). Regional disparities observed here can, however, be linked to specific circumstances; for example, the lower mean OGP found in the Mediterranean is likely related to growth limitation through lower food availability in this largely oligotrophic region (Siokou-Frangou et al., 2010), a view supported by laboratory experiments on bryozoans (Svensson & Marshall, 2015) and Antarctic bivalves (Román-González et al., 2017). Such observations emphasise that, whilst deviation from the global mean can form an important means of determining local effects that are driven by subtle ecological variation, spatial analysis of a larger biogeographical range can highlight variation in physiological characteristics otherwise invisible in studies conducted over smaller scales (Chown, Gaston, & Robinson, 2004; Parmesan & Yohe, 2003). This is important as unique local environmental characteristics and history may have a greater affect than previously recognised when data has been pooled from perceived identical environments.

Whilst our analysis confirms that overall growth performance at lower temperature deviates from the expected thermodynamic relationship (e.g. Peck, 2016), we identify a critical difference between

Arctic and Antarctic environments that has not been previously recognised or distinguished (Heilmayer et al., 2004; Peck, 2016). Specifically, when average bottom temperatures are  $< 0^{\circ}\text{C}$ , the growth of Arctic bivalve species are consistent with physiological expectations, whereas the growth of Antarctic bivalve species deviates below the expected relationship. The only exception to this pattern is the Antarctic scallop *Adamussium colbecki* which remains within expectation. This revelation is striking, because *A. colbecki* is routinely used as a model species for physiological studies in the Antarctic (Moro et al., 2019), meaning that *a priori* assessments of the physiological responses of polar ecosystems to climate warming may underestimate species vulnerability in the Antarctic whilst overestimating species vulnerability in the Arctic. Further, differences in growth performance between subclasses are likely to be reflected in their overall physiology (Pörtner et al., 2001); the subclass Pteriomorphia contains fewer, but larger, families, which include commercially valuable species (van der Schatte Olivier et al., 2018), such as Mussels (Mytilidae), Oysters (Ostreidae), and Scallops (Pectinidae), which may dominate regional assessments of physiological fitness. Should a high proportion of studies focus on a limited subset of regionally adapted species, or physiological typology, there is potential for model projections to perpetuate skewed conclusions about the most likely effects of climate change at larger scales (Wernberg, Smale, & Thomsen, 2012). These observations are consistent with other research in global marine environments that has shown regional and temporal differentiation in the severity and direction of effects associated with climatic forcing (Dijkstra, Westerman, & Harris, 2011; Godbold & Solan, 2013).

Although our analyses confirm a positive relationship between sea temperatures and decreasing latitude, it is important to emphasise that sea temperature is also influenced by local environmental cycles and other phenomena (e.g. depth, upwelling, El Niño) which can affect regional physiological responses. In this respect, it is noteworthy that the large degree of overlap of the Arctic and Temperate latitudinal zone reflects the boreal origins of many benthic species (Piepenburg, 2005), and that the large degree of overlap in high POC in the Arctic and Temperate regions reflects food availability to the benthos and species distribution (Solan et al. 2020). A key characteristic of the Arctic is the Atlantic influence and overlap of species distributions and physiological responses over the polar front (Piepenburg, 2005; Richard, Morley, Deloffre, & Peck, 2012), which contrasts to the Antarctic which is effectively isolated by the Antarctic Circumpolar Current from the southern Temperate Zone (Clarke & Crame, 2010) resulting in a relatively long evolutionary isolation of the Southern Ocean (Chown et al., 2015; Crame et al., 2014; Clarke, Barnes, & Hodgson, 2005). Isolation over evolutionary relevant timescales in the Antarctic has led to unique fauna and adaptations in response to low temperature (Barnes, Fuentes, Clarke, Schloss, & Wallace, 2006) which slow their biological processes beyond expectations (Peck, 2016; Peck, Heiser, & Clark, 2016). Hence, we observe greater thermal sensitivity and reduced OGP in the Antarctic relative to the Arctic (Richard et al., 2012). An important implication of this finding is that the common practice of pooling data from realms that are perceived as being similar to one another in order to overcome paucity of data is undesirable, as species responses are unlikely to be uniform across ecoregions.

The activation energies, representing all the reactions involved in the synthesis of proteins for growth, and calcification of the shell, strongly indicate that tropical realms and the Southern Ocean will be disproportionately affected by projected climate change in the 21<sup>st</sup> Century (IPCC, 2018), albeit with different population responses. A decrease in growth rate with increasing temperatures suggests that species may be beyond their thermal optimum, and is supported by theories of tropical species living close to their thermal limits (Dell, Pawer, & Savage, 2011; Amarasekare and Savage, 2012). Despite the importance of temperature however, the maximum rates of growth in an individual is unlikely to be achieved because of, for example, resource limitation (Siokou-Frangou et al., 2010) or highly seasonal food input (Zuo, Moses, West, Hou, & Brown, 2012). The regions of negative or very high activation energy in our study reflect these two environmental conditions. In the Antarctic, where temperature variations are very small, seasonality in metabolism can positively relate to food input in species of sponge (Dayton, Robilliard, Paine, & Dayton, 1974), echinoderm (Brockington & Clarke, 2001), and bryozoan (Barnes, 1995), which show faster growth and metabolic rates than expected during periods of high food availability. However, immature Southern Ocean *A. colbecki* show no uncoupling of metabolic rate from temperature, suggesting an ontogenetic component to relationships between somatic growth, food availability, and temperature (Heilmayer et al., 2005). The Arctic is also food limited with an observed mismatch between shell growth and body mass in the bivalves *Serripes groenlandicus* and *Chlamys islandica* before the onset of phytoplankton bloom (Blicher, Rysgaard, & Sejr, 2010; Carroll et al., 2011). Here, food quality rather than quantity is shown to be fundamental factor, and can be observed through the transition of lower quality phytoplankton to nutrient rich sea ice algae during the seasonal sea ice retreat, and a projected negative response to warming related changes in sea ice primary production in Arctic fjords (Carroll et al., 2011).

An alternative explanation to the constrained growth in Antarctic bivalves is protein synthesis, a vital process in somatic growth widely hypothesised to be limited at low or high temperatures (Dell, Pawer, & Savage, 2011; Peck, 2016). How this might be negated at low temperatures by projected warming is, however, currently unclear (Clark et al., 2019). The pathways involved in the synthesis and folding of functional proteins, generally occurs faster as temperatures increase and is widely postulated to be expensive and sensitive at low temperature (Clarke, 2017; Fraser & Rogers, 2007). However, adaptive changes within the genome can overcome protein efficiency to some extent (Chen et al., 2008). The sensitivities of growth performance to warming in the tropical and polar biogeographical realms may reflect these molecular constraints, however, research into protein function and synthesis in marine organisms is still in its infancy (Tomanek, 2011). This emerging area of research may well elucidate the molecular mechanisms behind limitations to growth in the lowest and highest temperature environments, and identify the constraints that maintain their sensitivity to temperature changes (Clark et al., 2019), especially when comparing the contrasting physiological responses from the Arctic and Antarctic realms from an evolutionary perspective (Feder, Bennett, & Huey, 2000).

Macroecological approaches to examine growth rates of important marine ectotherms, as used here, form an essential link between laboratory experiments and appropriate regionally adjusted assessment of risk (Chown, Gaston, et al., 2004; Chown, Sinclair, et al., 2004). The non-linear

relationship of OGP with temperature and latitude suggests that other ecological and phylogenetic constraints will exist across a latitudinal gradient and between ecological realms (Parmesan & Yohe, 2003), but these may be underestimated or ignored when traditional assumptions on thermal relationships are applied in isolation. This carries implications to the way we assess physiological responses to climate change scenarios (Clarke, 2003; Feder et al., 2000) as the consequences of climate change will differ at local scales (Stuart-Smith et al., 2015), or across species distributions (Deutsch, Penn, & Siebel, 2020). Climate change is not consistently expressed across latitude ranges and while experimental approaches have persistently shown the greatest severity of warming on physiology to be within tropical (Deutsch et al., 2008; Tewksbury, Huey, & Deutsch, 2008) and across both polar realms (Peck et al., 2004; Pörtner, Peck, & Somero, 2007), contradictory linear relationships with greatest thermal capacity at the tropics have also been identified (Seebacher et al., 2014). With this in mind, identification of where management and conservation efforts should be focussed can be achieved by using better fitting statistical models, which do not assume linearity in thermal relationships with physiological parameters. Linking regional and global physiological patterns with acclimation capacity, plasticity, and ultimately adaptation to projected environmental change will be essential to ameliorating the consequences of climate change on ecosystem function to protect against the loss of ecosystem services.

**Data Accessibility**

The growth and corresponding environmental data (Particulate Organic Carbon and Temperature) used in this study is available through figshare with appropriate descriptors (<https://doi.org/10.6084/m9.figshare.9943058.v1>)

**References**

Amante, C. and B.W. Eakins, 2009. ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. *National Geophysical Data Center*, NOAA. doi:10.7289/V5C8276

Amarasekare, P., and Savage, V. (2012) A framework for elucidating the temperature dependence of fitness, *American Society of Naturalists*, 179, 178 – 191. doi: 10.1086/663677

Barnes, D. K. A. (1995). Seasonal and annual growth in erect species of Antarctic Bryozoans *Journal of Experimental Marine Biology and Ecology*, 188(2), 181-198. doi:10.1016/0022-0981(95)00003-a

Barnes, D. K. A., Fuentes, V., Clarke, A., Schloss, I. R., & Wallace, M. I. (2006). Spatial and temporal variation in shallow seawater temperatures around Antarctica. *Deep-Sea Research Part II: Topical Studies in Oceanography*, 53(8-10), 853-865. doi:10.1016/j.dsr2.2006.03.008

Berkman, P. A., Cattaneo-Vietti, R., Chiantore, M., & Howard-Williams, C. (2004). Polar emergence and the influence of increased sea-ice extent on the Cenozoic biogeography of pectinid molluscs in Antarctic coastal areas. *Deep Sea Research Part II: Topical Studies in Oceanography*, 51(14), 1839-1855. doi:<https://doi.org/10.1016/j.dsr2.2004.07.017>

Blicher, M. E., Rysgaard, S., & Sej, M. K. (2010). Seasonal growth variation in *Chlamys islandica* (Bivalvia) from sub-Arctic Greenland is linked to food availability and temperature. *Marine Ecology Progress Series*, 407, 71-86. doi:10.3354/meps08536

Brey, T. (1999). Growth performance and mortality in aquatic macrobenthic invertebrates. *Advances in Marine Biology*, 35, 153-223.

Brierley, A. S., & Kingsford, M. J. (2009). Impacts of Climate Change on Marine Organisms and Ecosystems. *Current Biology*, 19(14), R602-R614. doi:10.1016/j.cub.2009.05.046



- Brockington, S., & Clarke, A. (2001). The relative influence of temperature and food on the metabolism of a marine invertebrate. *Journal of Experimental Marine Biology and Ecology*, 258(1), 87-99. doi:10.1016/S0022-0981(00)00347-6
- Burrows, M. T., Schoeman, D. S., Buckley, L. B., Moore, P., Poloczanska, E. S., Brander, K. M., . . . Richardson, A. J. (2011). The Pace of Shifting Climate in Marine and Terrestrial Ecosystems. *Science*, 334(6056), 652-655. doi:10.1126/science.1210288
- Calosi, P., De Wit, P., Thor, P., & Dupont, S. (2016). Will life find a way? Evolution of marine species under global change. *Evolutionary Applications*, 9(9), 1035-1042. doi:10.1111/eva.12418
- Carr, M.-E. (2001) Estimation of potential productivity in Eastern Boundary Currents using remote sensing, Deep Sea Research., Part II, 49(1-3), 59-80, doi:[10.1016/S0967-0645\(01\)00094-7](https://doi.org/10.1016/S0967-0645(01)00094-7)
- Carroll, M. L., Ambrose, W. G., Levin, B. S., Locke, W. L., Henkes, G. A., Hop, H., & Renaud, P. E. (2011). Pan-Svalbard growth rate variability and environmental regulation in the Arctic bivalve *Serripes groenlandicus*. *Journal of Marine Systems*, 88(2), 239-251. doi:10.1016/j.jmarsys.2011.04.010
- Chen, Z., Cheng, C.-H. C., Zhang, J., Cao, L., Chen, L., Zhou, L., . . . Chen, L. (2008). Transcriptomic and genomic evolution under constant cold in Antarctic notothenioid fish. *Proceedings of the National Academy of Sciences*, 105(35), 12944-12949. doi:10.1073/pnas.0802432105
- Chown, S. L., Clarke, A., Fraser, C. I., Cary, S. C., Moon, K. L., & McGeoch, M. A. (2015). The changing form of Antarctic biodiversity. *Nature*, 522(7557), 431.
- Chown, S. L., Gaston, K. J., & Robinson, D. (2004). Macrophysiology: large-scale patterns in physiological traits and their ecological implications. *Functional Ecology*, 18(2), 159-167. doi:10.1111/j.0269-8463.2004.00825.x
- Chown, S. L., Sinclair, B. J., Leinaas, H. P., & Gaston, K. J. (2004). Hemispheric asymmetries in biodiversity - A serious matter for ecology. *PLOS Biology*, 2(11), 1701-1707. doi:10.1371/journal.pbio.0020406
- Clark, M. S., Suckling, C. C., Cavallo, A., Mackenzie, C. L., Thorne, M. A. S., Davies, A. J., & Peck, L. S. (2019). Molecular mechanisms underpinning transgenerational plasticity in the green sea urchin *Psammechinus miliaris*. *Scientific Reports*, 9(1), 952. doi:10.1038/s41598-018-37255-6
- Clarke, A. (2003). Costs and consequences of evolutionary temperature adaptation. *Trends in Ecology & Evolution*, 18(11), 573-581. doi:10.1016/j.tree.2003.08.007
- Clarke, A. (2017). *Principles of thermal ecology: temperature, energy and life*: Oxford University Press.
- Clarke, A., Barnes, D. K., & Hodgson, D. A. (2005). How isolated is Antarctica? *Trends in Ecology & Evolution*, 20(1), 1-3.
- Clarke, A., & Crame, J. A. (2010). Evolutionary dynamics at high latitudes: speciation and extinction in polar marine faunas. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 365(1558), 3655-3666. doi:10.1098/rstb.2010.0270
- Clarke, A., & Johnston, N. M. (2003). Antarctic marine benthic diversity. *Oceanography and Marine Biology*, Vol 41, 41, 47-114.
- Crame, J. A., Beu, A. G., Ineson, J. R., Francis, J. E., Whittle, R. J., & Bowman, V. C. (2014). The Early Origin of the Antarctic Marine Fauna and Its Evolutionary Implications. *Plos One*, 9(12), e114743. doi:10.1371/journal.pone.0114743
- Dayton, P., Jarrell, S., Kim, S., Thrush, S., Hammerstrom, K., Slattery, M., & Parnell, E. (2016). Surprising episodic recruitment and growth of Antarctic sponges: Implications for ecological resilience. *Journal of Experimental Marine Biology and Ecology*, 482, 38-55. doi:10.1016/j.jembe.2016.05.001
- Dayton, P. K., Robilliard, G. A., Paine, R. T., & Dayton, L. B. (1974). Biological Accommodation in the Benthic Community at McMurdo Sound, Antarctica. *Ecological Monographs*, 44(1), 105-128. doi:10.2307/1942321
- Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy of Sciences*, 108(26), 10591-10596.
- Dell'Acqua, O., Trębala, M., Chiantore, M., & Hannula, S.-P. (2019). Robustness of *Adamussium colbecki* shell to ocean acidification in a short-term exposure. *Marine Environmental Research*, 149, 90-99. doi:<https://doi.org/10.1016/j.marenvres.2019.06.010>
- Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, 105(18), 6668-6672. doi:10.1073/pnas.0709472105

- Dijkstra, J. A., Westerman, E. L., & Harris, L. G. (2011). The effects of climate change on species composition, succession and phenology: a case study. *Global Change Biology*, 17(7), 2360-2369. doi:10.1111/j.1365-2486.2010.02371.x
- Feder, M. E., Bennett, A. F., & Huey, R. B. (2000). Evolutionary Physiology. *Annual Review of Ecology and Systematics*, 31(1), 315-341. doi:10.1146/annurev.ecolsys.31.1.315
- Fraser, K. P. P., & Rogers, A. D. (2007). Protein Metabolism in Marine Animals: The Underlying Mechanism of Growth. *Advances in Marine Biology*, 52, 267-362. doi:10.1016/S0065-2881(06)52003-6
- Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., & Charnov, E. L. (2001). Effects of Size and Temperature on Metabolic Rate. *Science*, 293(5538), 2248.
- Gladstone-Gallagher, R. V., Pilditch, C. A., Stephenson, F., & Thrush, S. F. (2019). Linking Traits across Ecological Scales Determines Functional Resilience. *Trends in Ecology & Evolution*. doi:10.1016/j.tree.2019.07.010
- Godbold, J. A., & Solan, M. (2013). Long-term effects of warming and ocean acidification are modified by seasonal variation in species responses and environmental conditions. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 368(1627). doi:10.1098/rstb.2013.0186
- Heilmayer, O., Brey, T., & Pörtner, H. O. (2004). Growth efficiency and temperature in scallops: a comparative analysis of species adapted to different temperatures. *Functional Ecology*, 18(5), 641-647.
- Heilmayer, O., Honnen, C., Jacob, U., Chiantore, M., Cattaneo-Vietti, R., & Brey, T. (2005). Temperature effects on summer growth rates in the Antarctic scallop, *Adamussium colbecki*. *Polar Biology*, 28(7), 523-527. doi:10.1007/s00300-005-0716-7
- Henson, S., R. Sanders, E. Madsen, P. Morris, F. Le Moigne and G. Quartly (2011), A reduced estimate of the strength of the ocean's biological carbon pump, *Geophysical Research Letters*, 38, L04606, doi:10.1029/2011GL046735
- Henson, S., R. Sanders and E. Madsen (2012), Global patterns in efficiency of particulate organic carbon export and transfer to the deep ocean, *Global Biogeochemical Cycles*, 26, GB1028, doi:10.1029/2011GB004099
- Hoegh-Guldberg, O., & Bruno, J. F. (2010). The Impact of Climate Change on the World's Marine Ecosystems. *Science*, 328(5985), 1523-1528. doi:10.1126/science.1189930
- IPCC. (2018). *Global Warming of 1.5° C: An IPCC Special Report on the Impacts of Global Warming of 1.5° C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*: Intergovernmental Panel on Climate Change.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., O. K. Baranova, . . . Seidov, D. (2013). *World Ocean Atlas 2013, Volume 1: Temperature* (Vol. 73): NOAA Atlas NESDIS.
- Martin, J. H., G. A. Knauer, D. M. Karl, and W. W. Broenkow (1987), VERTEX: Carbon cycling in the Northeast Pacific, *Deep Sea Research*, Part A, 34(2), 267-285, doi:10.1016/0198-0149(87)90086-0
- Moro, G., Buonocore, F., Barucca, M., Spazzali, F., Canapa, A., Pallavicini, A., . . . Gerdol, M. (2019). The first transcriptomic resource for the Antarctic scallop *Adamussium colbecki*. *Marine Genomics*, 44, 61-64. doi:10.1016/j.margen.2018.09.007
- Moss, D. K., Ivany, L. C., Judd, E. J., Cummings, P. W., Bearden, C. E., Kim, W. J., . . . Driscoll, J. R. (2016). Lifespan, growth rate, and body size across latitude in marine Bivalvia, with implications for Phanerozoic evolution. *Proceedings of the Royal Society B-Biological Sciences*, 283(1836), 7. doi:10.1098/rspb.2016.1364
- Moss, D. K., Ivany, L. C., Silver, R. B., Schue, J., & Artruc, E. G. (2017). High-latitude settings promote extreme longevity in fossil marine bivalves. *Paleobiology*, 43(3), 365-382. doi:10.1017/pab.2017.5
- Ohlberger, J. (2013). Climate warming and ectotherm body size – from individual physiology to community ecology. *Functional Ecology*, 27(4), 991-1001. doi:10.1111/1365-2435.12098
- Osovitz, C. J., & Hofmann, G. E. (2007). Marine macrophysiology: Studying physiological variation across large spatial scales in marine systems. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 147(4), 821-827. doi:10.1016/j.cbpa.2007.02.012
- Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421(6918), 37.
- Pebesma, E., Graeler, B., & Pebesma, M. E. (2019). Package 'gstat'.



- Peck, L. S. (2016). A Cold Limit to Adaptation in the Sea. *Trends in Ecology & Evolution*, 31(1), 13-26.
- Peck, L. S. (2018). *Antarctic Marine Biodiversity: Adaptations, Environments and Responses to Change: An Annual Review*.
- Peck, L. S., Heiser, S., & Clark, M. S. (2016). Very slow embryonic and larval development in the Antarctic limpet *Nacella polaris*. *Polar Biology*, 39(12), 2273-2280. doi:10.1007/s00300-016-1894-1
- Peck, L. S., Webb, K. E., & Bailey, D. M. (2004). Extreme sensitivity of biological function to temperature in Antarctic marine species. *Functional Ecology*, 18(5), 625-630. doi:10.1111/j.0269-8463.2004.00903.x
- Piepenburg, D. (2005). Recent research on Arctic benthos: common notions need to be revised. *Polar Biology*, 28(10), 733-755. doi:10.1007/s00300-005-0013-5
- Pinheiro, J., & Bates, D. (2000). *Mixed-effects models in S and S-PLUS*. New York, LCC: Springer Verlag.
- Pinheiro, J., Bates, D., DebRoy, S., & Sarkar, D. (2018). R Core Team (2018). nlme: Linear and nonlinear mixed effects models. R package version 3.1-124. In.
- Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, 569(7754), 108-111. doi:10.1038/s41586-019-1132-4
- Pörtner, H. O. (2001). Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften*, 88(4), 137-146.
- Pörtner, H. O., Berdal, B., Blust, R., Brix, O., Colosimo, A., De Wachter, B., . . . Zakhartsev, M. (2001). Climate induced temperature effects on growth performance, fecundity and recruitment in marine fish: developing a hypothesis for cause and effect relationships in Atlantic cod (*Gadus morhua*) and common eelpout (*Zoarces viviparus*). *Continental Shelf Research*, 21(18-19), 1975-1997.
- Pörtner, H.O., Storch, D., & Heilmayer O. (2005). Constraints and trade-offs in climate dependant adaptation: energy budgets and growth in a latitudinal cline. *Scientia Marina*, 69, 271 - 285
- Pörtner, H. O., & Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science*, 315(5808), 95-97.
- Pörtner, H. O., Peck, L., & Somero, G. (2007). Thermal limits and adaptation in marine Antarctic ectotherms: an integrative view. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 362(1488), 2233-2258. doi:10.1098/rstb.2006.1947
- Reed, A. J., Linse, K., & Thatje, S. (2014). Differential adaptations between cold-stenothermal environments in the bivalve *Lissarca cf. miliaris* (Philobryidae) from the Scotia Sea islands and Antarctic Peninsula. *Journal of Sea Research*, 88, 11-20.
- Reed, A. J., Godbold, J. A., Grange, L.J., Solan, M. (2020): A global database of marine bivalve growth rates and parameters derived from von bertalanffy growth curves. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.9943058.v1>
- Richard, J., Morley, S. A., Deloffre, J., & Peck, L. S. (2012). Thermal acclimation capacity for four Arctic marine benthic species. *Journal of Experimental Marine Biology and Ecology*, 424, 38-43. doi:10.1016/j.jembe.2012.01.010
- Román-González, A., Scourse, J. D., Butler, P. G., Reynolds, D. J., Richardson, C. A., Peck, L. S., . . . Hall, I. R. (2017). Analysis of ontogenetic growth trends in two marine Antarctic bivalves *Yoldia eightsi* and *Laternula elliptica*: Implications for sclerochronology. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 465, 300-306. doi:10.1016/j.palaeo.2016.05.004
- Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C., & Pounds, J. A. (2003). Fingerprints of global warming on wild animals and plants. *Nature*, 421(6918), 57.
- Seebacher, F., White, C. R., & Franklin, C. E. (2014). Physiological plasticity increases resilience of ectothermic animals to climate change. *Nature Climate Change*, 5, 61. doi:10.1038/nclimate2457
- Seneviratne, S. I., Rogelj, J., Séférian, R., Wartenburger, R., Allen, M. R., Cain, M., . . . Warren, R. F. (2018). The many possible climates from the Paris Agreement's aim of 1.5 °C warming. *Nature*, 558(7708), 41-49. doi:10.1038/s41586-018-0181-4
- Siokou-Frangou, I., Christaki, U., Mazzocchi, M. G., Montresor, M., d'Alcala, M. R., Vaque, D., & Zingone, A. (2010). Plankton in the open Mediterranean Sea: a review. *Biogeosciences*, 7(5), 1543-1586. doi:10.5194/bg-7-1543-2010
- Solan, M., Ward, E.R., Wood, C.L., Reed, A.J., Grange, L.J. and Godbold, J.A. (2020). Climate-driven benthic invertebrate activity and biogeochemical functioning across the Barents Sea polar front. *Philosophical Transactions of the Royal Society*, 378:20190365. doi/10.1098/rsta.2019.0365

1  
2  
3 604 Somero, G. N. (2010). The physiology of climate change: how potentials for acclimatization and  
4 605 genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*,  
5 606 213(6), 912-920. doi:10.1242/Jeb.037473  
6 607 Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdaña, Z. A., Finlayson, M., . . . Lourie, S. A.  
7 608 (2007). Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *AIBS*  
8 609 *Bulletin*, 57(7), 573-583.  
9 610 Stuart-Smith, R. D., Edgar, G. J., Barrett, N. S., Kininmonth, S. J., & Bates, A. E. (2015). Thermal  
10 611 biases and vulnerability to warming in the world's marine fauna. *Nature*, 528(7580), 88.  
11 612 doi:10.1038/nature16144  
12 613 Svensson, J. R., & Marshall, D. J. (2015). Limiting resources in sessile systems: food enhances  
13 614 diversity and growth of suspension feeders despite available space. *Ecology*, 96(3), 819-827.  
14 615 doi:10.1890/14-0665.1  
15 616 Team, R. C. (2018). R: A language and environment for statistical computing. Vienna, Austria.  
16 617 Retrieved from <https://www.R-project.org/>  
17 618 Tewksbury, J. J., Huey, R. B., & Deutsch, C. A. (2008). Putting the Heat on Tropical Animals.  
18 619 *Science*, 320(5881), 1296. doi:10.1126/science.1159328  
19 620 Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., . . .  
20 621 Hannah, L. (2004). Extinction risk from climate change. *Nature*, 427(6970), 145.  
21 622 Tomanek, L. (2011). Environmental Proteomics: Changes in the Proteome of Marine Organisms in  
22 623 Response to Environmental Stress, Pollutants, Infection, Symbiosis, and Development.  
23 624 *Annual Review of Marine Science*, 3(1), 373-399. doi:10.1146/annurev-marine-120709-  
24 625 142729  
25 626 Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, 348(6234), 571-573.  
26 627 doi:10.1126/science.aaa4984  
27 628 van der Schatte Olivier, A., Jones, L., Vay, L. L., Christie, M., Wilson, J., & Malham, S. K. (2018). A  
28 629 global review of the ecosystem services provided by bivalve aquaculture. *Reviews in*  
29 630 *Aquaculture*, 0(0). doi:10.1111/raq.12301  
30 631 Watson, S. A., Morley, S. A., & Peck, L. S. (2017). Latitudinal trends in shell production cost from the  
31 632 tropics to the poles. *Science Advances*, 3(9), 8. doi:10.1126/sciadv.1701362  
32 633 Wernberg, T., Smale, D. A., & Thomsen, M. S. (2012). A decade of climate change experiments on  
33 634 marine organisms: procedures, patterns and problems. *Global Change Biology*, 18(5), 1491-  
34 635 1498. doi:10.1111/j.1365-2486.2012.02656.x  
35 636 Wohlgemuth, D., Solan, M., & Godbold, J. A. (2016). Specific arrangements of species dominance  
36 637 can be more influential than evenness in maintaining ecosystem process and function.  
37 638 *Scientific Reports*, 6, 39325. doi:10.1038/srep39325  
38 639 Wood, S. (2019). Package 'mgcv': Mixed GAM Computation Vehicle with Automatic Smoothness  
39 640 Estimation. R Package version 1.8 - 29.  
40 641 WoRMS Editorial Board (2020). World Register of Marine Species. Available from  
41 642 <http://www.marinespecies.org> at VLIZ. Accessed 2019-09-01. doi:10.14284/170  
42 643 Yamahira, K., & Conover, D. O. (2002). Intra- vs. Interspecific latitudinal variation in growth:  
43 644 Adaptation to temperature or seasonality? *Ecology*, 83(5), 1252-1262. doi:10.1890/0012-  
44 645 9658(2002)083  
45 646 Zuo, W., Moses, M. E., West, G. B., Hou, C., & Brown, J. H. (2012). A general model for effects of  
46 647 temperature on ectotherm ontogenetic growth and development. *Proceedings of the Royal*  
47 648 *Society B: Biological Sciences*, 279(1734), 1840-1846. doi:10.1098/rspb.2011.2000  
48 649 Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and*  
49 650 *extensions in ecology with R*: Springer Verlag, New York, LLC.  
50 651 Zuur, A., Saveliev, A.A., Ieno, E.N. (2015) *A Beginner's Guide to Generalised Additive Mixed Models*  
51 652 *with R*. Highland Statistics Ltd.

52 653  
53 654  
54 655  
55  
56 656 **Figure Legends**

57 657 **Figure 1. Variation in overall growth performance with latitude, realm, and biogeographic**  
58 658 **province** a) Trend in overall growth performance with latitude, b) Mean overall growth performance ( $\pm$   
59 659 95% confidence intervals) in each biogeographical Realm, and c) Global map of mean overall growth

performance within each biogeographic Province and data points overlaid with colour representing mean annual average bottom temperature. In a) Model prediction (solid line) and 95% confidence intervals (dotted line) for changes in overall growth performance with latitude are shown, with data point colours representing the four latitudinal zones.


**Figure 2. Trends in overall growth performance of bivalves with a) average annual bottom temperature (°C) and b) particulate organic carbon (ln g C m<sup>-2</sup> yr<sup>-1</sup>) and in c) overall growth performance for each Subclass.** In a) and b) model predictions (solid lines) and 95% confidence intervals (dashed lines) are shown for changes in overall growth performance for the Subclasses Heterodonta (light blue) and Pteriomorphia (light blue); c) median is indicated at the midpoint, the upper and lower quartiles are indicated by the hinges, and open circles indicate outliers. Data points are superimposed.

**Figure 3. Geographically differentiated relationships of overall growth performance and absolute temperature** a) Arrhenius model of global overall growth performance values against inverse temperature (in Kelvins) (n.b. temperature scale from high to low); b) Arrhenius model using Arctic (dark blue) and Antarctic (red) data points with regression line of the global relation of overall growth performance to inverse temperature (in Kelvins).


**Figure 4. Activation energies derived from Arrhenius models for geographic regions** a) Activation energy (ev) as calculated from the slope of the line in each latitude zone; b) Activation energy (ev) as calculated from the slope of the line within each geographical Realm. Realm codes given in parenthesis; 1 – Arctic; 2 – Temperate North Atlantic; 3 – Temperate North Pacific; 4 – Tropical Atlantic; 5 – West Indo-Pacific; 6 – Central Indo-Pacific; 7 – East Indo-Pacific; 8 – Tropical East Pacific; 9 – Temperate South America; 10 – Temperate South Africa; 11 – Temperate Australia; 12 – Southern Ocean.

Figure 1

a



b



c






Figure 2




Figure 3





Figure 4





## References relating to the meta-analysis data set.

- Abada-Boudjema, Y., Altes, J., & Moueza, M. (1984). Growth of two species of mussels, *Mytilus galloprovincialis* and *Perna perna* in a natural mussel bed in the Bay of Algiers. *Haliotis*, 14, 33-38.
- Abele, D., Strahl, J., Brey, T., & Philipp, E. E. R. (2008). Imperceptible senescence: Ageing in the ocean quahog *Arctica islandica*. *Free Radical Research*, 42(5), 474-480.
- Abrahao, J. R., Cardoso, R. S., Yokoyama, L. Q., & Amaral, A. C. Z. (2010). Population biology and secondary production of the stout razor clam *Tagelus plebeius* (Bivalvia, Solecurtidae) on a sandflat in southeastern Brazil. *Zoologia*, 27(1), 54-64. doi:10.1590/s1984-46702010000100009
- Acarli, S., Lok, A., & Yigitkurt, S. (2012). Growth and survival of *Anadara inaequalis* (Bruguiere, 1789) in Sufa Lagoon, Izmir, Turkey. *The Israeli Journal of Aquaculture - Bamidgeh*, 64, 7.
- Al-Barwani, S. M., Arshad, A., Amin, S. M. N., Japar, S. B., Siraj, S. S., & Yap, C. K. (2007). Population dynamics of the green mussel *Perna viridis* from the high spat-fall coastal water of Malacca, Peninsular Malaysia. *Fisheries Research*, 84(2), 147-152. doi:https://doi.org/10.1016/j.fishres.2006.10.021
- Alagaraswami, K. (1966). Studies on some aspects of biology of the wedge-clam, *Donax faba* Gmelin from Mandapam coast in the Gulf of Mannar. *Journal of the Marine Biological Association of India*, 8, 56-75.
- Allison, E. (1994). Seasonal growth models for great scallops (*Pecten maximus* (L.)) and queen scallops (*Aequipecten opercularis* (L.)). *Journal of Shellfish Research*, 13(2), 555-564.
- Allison, E. H., Wilson, U. A. W., & Brand, A. R. (1994). Age determination and the first growth ring in North Irish Sea populations of the scallop, *Pecten maximus* (L.). *Journal of Molluscan Studies*, 60(1), 91-95. doi:10.1093/mollus/60.1.91
- Ambrose, W. G., Carroll, M. L., Greenacre, M., Thorrold, S. R., & McMahon, K. W. (2006). Variation in *Serripes groenlandicus* (Bivalvia) growth in a Norwegian high-Arctic fjord: evidence for local- and large-scale climatic forcing. *Global Change Biology*, 12(9), 1595-1607. doi:10.1111/j.1365-2486.2006.01181.x
- Ambrose, W. G., Renaud, P. E., Locke, W. L., Cottier, F. R., Berge, J., Carroll, M. L., . . . Ryan, S. (2012). Growth line deposition and variability in growth of two circumpolar bivalves (*Serripes groenlandicus*, and *Clinocardium ciliatum*). *Polar Biology*, 35(3), 345-354.
- Amin, H., Tenjing, Y., & Thippeswamy, S. (2017). Population dynamics of the Asian green mussel *Perna viridis* (L.) from St. Mary's islands off Malpe, India. *Indian Journal of Geo-Marine Sciences*, 46, 1659-1666.
- Ansel, A. (1961). Reproduction, growth and mortality of (*Venus striatula*) in Kame Bay, Millport. *Journal of the Marine Biological Association of the United Kingdom*, 41, 191-215.
- Ansell, A. D., & Lagardère, F. (1980). Observations on the biology of *Donax trunculus* and *D. vittatus* at Ile d'Oléron (French Atlantic coast). *Marine Biology*, 57(4), 287-300. doi:10.1007/bf00387572
- Ansell, A. D., Sivadas, P., Narayanan, B., Sankaranarayanan, V. N., & Trevallion, A. (1972). The ecology of two sandy beaches in south west India. I. Seasonal changes in physical and chemical factors, and in the macrofauna. *Marine Biology*, 17(1), 38-62. doi:10.1007/bf00346953
- Antoine, L. (1979). La croissance de la coquille Saint-Jacques *Pecten maximus* (L.) et ses variations en mer Celtique et en Manche. (PhD thesis), University of Bretagne Occidentale, Brest. 148 pp.
- Anwar, N. A., Richardson, C. A., & Seed, R. (1990). Age determination, growth rate and population structure of the horse mussel *Modiolus modiolus*. *Journal of the Marine Biological Association of the United Kingdom*, 70(2), 441-457. doi:10.1017/S0025315400035529
- Appeldoorn, R. S. (1995). Covariation in life-history parameters of soft-shell clams (*Mya arenaria*) along a latitudinal gradient. *ICES Marine Science Symposium*, 199, 12-25.

- Appukuttan, K. K. (1999). Population dynamics of an exploited stock of the clam *Paphia malabarica* of Ashtamudi Estuary. *The Fourth Indian Fisheries Forum Proceedings*, 31-34.
- Aragon-Noriega, E. A. (2013). Individual growth modeling of the penshell *Atrina maura* (Bivalvia: Pinnidae) using a multi model inference approach. *Revista De Biología Tropical*, 61(3), 1167-1174.
- Aravindakshan, I. (1955). *Studies on the biology of the queen scallop, Chlamys opercularis (L.)*. (Ph.D Thesis), University of Leeds, UK.
- Argente, F. A., & Estacion, J. (2014). *Effect of different harvesting practices on the dynamics of Paphia textile (Gmelin 1792) (Bivalvia: Veneridae) populations at two sites in Zamboanga del Norte, Southern Philippines* (Vol. 12).
- Arneri, E., Frogli, C., Polenta, R., & Antolini, B. (1997). Growth of *Chamelea gallina* (Bivalvia: Veneridae) in the eastern Adriatic (Neretva river estuary). *Tisucu Godina Prvoga Spomena Ribarstva u Hrvata*, 597, 669-676.
- Arneri, E., Giannetti, G., & Antolini, B. (1998). Age determination and growth of *Venus verrucosa* L. (Bivalvia: Veneridae) in the southern Adriatic and the Aegean Sea. *Fisheries Research*, 38(2), 193-198. doi:https://doi.org/10.1016/S0165-7836(98)00146-5
- Arneri, E., Giannetti, G., Polenta, R., & Antolini, B. (1995). Age and growth of *Chamelea gallina* (Bivalvia: Veneridae) in the Central Adriatic Sea obtained by thin sections. *Rapports et procès-verbaux des réunions Commission internationale pour l'exploration scientifique de la Mer Méditerranée*, 34, 17.
- Arnold, W. S., Marelli, D. C., Bert, T. M., Jones, D. S., & Quitmyer, I. R. (1991). Habitat-specific growth of hard clams *Mercenaria mercenaria* (L.) from the Indian River, Florida. *Journal of Experimental Marine Biology and Ecology*, 147(2), 245-265. doi:https://doi.org/10.1016/0022-0981(91)90185-Y
- Arntz, W. E., Brey, T., Tarazona, J., & Robles, A. (1987). Changes in the structure of a shallow sandy-beach community in Peru during an El Niño event. *South African Journal of Marine Science*, 5(1), 645-658. doi:10.2989/025776187784522504
- Arrieche, D., & Prieto, A. (2006). *Parámetros poblacionales del guacuco Tivela mactroides (Bivalvia: Veneridae) de Playa Caicara, Estado Anzoátegui, Venezuela* Population parameters of the trigonal tivala *Tivela mactroides* (Bivalvia: Veneridae) from Caicara Beach, Anzoátegui, Venezuela (Vol. 32).
- Bachelet, G. (1980). Growth and recruitment of the tellinid bivalve *Macoma balthica* at the southern limit of its geographical distribution, the Gironde estuary (SW France). *Marine Biology*, 59(2), 105-117. doi:10.1007/BF00405460
- Bagur, M., Richardson, C. A., Gutierrez, J. L., Arribas, L. P., Doldan, M. S., & Palomo, M. G. (2013). Age, growth and mortality in four populations of the boring bivalve *Lithophaga patagonica* from Argentina. *Journal of Sea Research*, 81, 49-56. doi:10.1016/j.seares.2013.04.003
- Barkai, A., & Branch, G. (1989). Growth and mortality of the mussels *Choromytilus meridionalis* (Krauss) and *Aulacomya ater* (Molina) as indicators of biotic conditions. *Journal of Molluscan Studies*, 55(3), 329-342. (Vol. 55).
- Barón, P. J., Real, L. E., Ciocco, N. F., & Ré, M. E. (2004). Morphometry, growth and reproduction of an Atlantic population of the razor clam *Ensis macha* (Molina, 1782). *Scientia Marina*, 68(2), 211-217.
- Barry, J. P., Whaling, P. J., & Kochevar, R. K. (2007). Growth, production, and mortality of the chemosynthetic vesicomyid bivalve, *Calyptogena kilmeri* from cold seeps off central California. *Marine Ecology-an Evolutionary Perspective*, 28(1), 169-182. doi:10.1111/j.1439-0485.2007.00119.x
- Bayne, B. L., & Worrall, C. M. (1980). Growth and Production of Mussels *Mytilus edulis* from two Populations. *Marine Ecology Progress Series*, 3(4), 317-328.
- Beaver, P. E., Bucher, D. J., & Joannes-Boyau, R. (2017). Growth patterns of three bivalve species targeted by the Ocean Cockle Fishery, southern New South Wales: *Eucrassatella kingicola*

- (Lamarck, 1805); *Glycymeris grayana* (Dunker, 1857); and *Callista (Notocallista) kingii* (Gray, 1827). *Journal of Molluscan Research*, 37(2), 104-112. doi:10.1080/13235818.2016.1253430
- Begum, S., Basova, L., Heilmayer, O., Philipp, E. E. R., Abele, D., & Brey, T. (2010). Growth and energy budget models of the bivalve *Arctica islandica* at six different sites in the Northeast Atlantic realm. *Journal of Shellfish Research*, 29(1), 107-115. doi:10.2983/035.029.0103
- Berg, C. J., & Alatalo, P. (1985). Biology of the tropical bivalve *Asaphis deflorata* (Linné, 1758). *Bulletin of Marine Science*, 37(3), 827-838.
- Berkman, P. (1990). The population biology of the Antarctic scallop, *Adamussium colbecki* (Smith 1902) at New Harbor, Ross Sea. In *Antarctic ecosystems* (pp. 281-288): Springer.
- Blicher, M. E., Rysgaard, S., & Sejr, M. K. (2010). Seasonal growth variation in *Chlamys islandica* (Bivalvia) from sub-Arctic Greenland is linked to food availability and temperature. *Marine Ecology Progress Series*, 407, 71-86. doi:10.3354/meps08536
- Blicher, M. E., Sejr, M. K., & Rysgaard, S. (2009). High carbon demand of dominant macrozoobenthic species indicates their central role in ecosystem carbon flow in a sub-Arctic fjord. *Marine Ecology Progress Series*, 383, 127-140. doi:10.3354/meps07978
- Boltacheva, N., & Mazlumyan, S. (2003). The growth and longevity of *Chamelea gallina* (Mollusca, Veneridae) in the Black Sea. *Vestnik Zoologii*, 37(3), 71-74.
- Boltachova, N., & Mazlumyan, S. (2001). The linear growth and lifetime of *Chamelea gallina* (Bivalvia: Veneridae) in the Black Sea. *Marine Ecology*, 55, 50-52.
- Bourne, N. (1982). Distribution, reproduction, and growth of Manila clam, *Tapes philippinarum* (Adams and Reeves), in British Columbia. *Journal of Shellfish Research*, 2, 47-54.
- Bourne, N., & Smith, D. (1972). *Breeding and growth of the horse clam, Tresus capax (Gould), in southern British Columbia*. Paper presented at the *Proceedings of the National Shellfisheries Association*.
- Boyden, C. (1972). Relationship of size to age in the cockles *Cerastoderma edule* and *C. glaucum* from the river Crouch estuary, Essex. *Journal of Conchology*, 27, 475-489.
- Breed-Willeke, G. M., & Hancock, D. (1980). Growth and reproduction of subtidal and intertidal populations of the gaper clam *Tresus capax* (Gould) from Yaquina Bay, Oregon. *Proceedings National Shellfisheries Association*, 70, 1-13.
- Breen, P. A., Gabriel, C., & Tyson, T. (1991). Preliminary estimates of age, mortality, growth, and reproduction in the hiatellid clam *Panopea zelandica* in New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 25(3), 231-237. doi:10.1080/00288330.1991.9516475
- Brêthes, J.-C. F., Desrosiers, G., & Fortin Jr, G. (1986). Croissance et production du bivalve *Mesodesma arctatum* (Conrad) sur la côte nord du golfe du Saint-Laurent. *Canadian Journal of Zoology*, 64(9), 1914-1919.
- Brey, T., Arntz, W., Pauly, D., & Rumohr, H. (1990). *Arctica (Cyprina) islandica* in Kiel Bay (Western Baltic): growth, production and ecological significance. *Journal of Experimental Marine Biology and Ecology*, 136(3), 217-235.
- Brey, T., & Clarke, A. (1993). Population-Dynamics of Marine Benthic Invertebrates in Antarctic and Sub-Antarctic Environments - Are There Unique Adaptations. *Antarctic Science*, 5(3), 253-266.
- Brey, T., & Hain, S. (1992). Growth, Reproduction and Production of *Lissarca notorcadensis* (Bivalvia, Philobryidae) in the Weddell Sea, Antarctica. *Marine Ecology Progress Series*, 82(3), 219-226.
- Briggs, R. P. (1982). Community Structure and Growth of *Mytilus edulis* L. in Lough Foyle. *Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and Chemical Science*, 82B, 245-259.
- Broom, M., & Mason, J. (1978). Growth and spawning in the pectinid *Chlamys opercularis* in relation to temperature and phytoplankton concentration. *Marine Biology*, 47(3), 277-285.
- Broom, M. J. (1982). Analysis of the Growth of *Anadara granosa* (Bivalvia, Arcidae) in Natural, Artificially Seeded and Experimental Populations. *Marine Ecology Progress Series*, 9(1), 69-79. doi:DOI 10.3354/meps009069

- 1
- 2
- 3 Brousseau, D. (1979). Analysis of growth rate in *Mya arenaria* using the Von Bertalanffy equation.
- 4 *Marine Biology*, 51(3), 221-227.
- 5 Brousseau, D., & Baglivo, J. (1987). A comparative study of age and growth in *Mya arenaria* (soft-
- 6 shell clam) from three populations in Long Island Sound. *Journal of Shellfish Research*, 6, 17-
- 7 24.
- 8 Brousseau, D. J. (1984). Age and Growth-Rate Determinations for the Atlantic Ribbed Mussel,
- 9 *Geukensia demissa*, Dillwyn (Bivalvia, Mytilidae). *Estuaries*, 7(3), 233-241. doi:Doi
- 10 10.2307/1352143
- 11 Brown, A., Heilmayer, O., & Thatje, S. (2010). Metabolic rate and growth in the temperate bivalve
- 12 *Mercenaria mercenaria* at a biogeographical limit, from the English Channel. *Journal of the*
- 13 *Marine Biological Association of the United Kingdom*, 90(5), 1019-1023.
- 14 doi:10.1017/s0025315409991470
- 15 Bureau, D. (2002). Age, size structure and growth parameters of geoducks (*Panopea abrupta*, Conrad
- 16 1849) from 34 locations in British Columbia sampled between 1993 and 2000. *Canadian*
- 17 *technical report of fisheries and aquatic sciences*, 2413, 1-84.
- 18 Bureau, D. (2003). Age, size structure and growth parameters of geoducks (*Panopea abrupta*, Conrad
- 19 1849) from seven locations in British Columbia sampled in 2001 and 2002. *Canadian*
- 20 *technical report of fisheries and aquatic sciences*, 2494, 1-29.
- 21 Bušelić, I., Peharda, M., Reynolds, D. J., Butler, P. G., González, A. R., Ezgeta-Balić, D., . . . Richardson,
- 22 C. A. (2015). *Glycymeris bimaculata* (Poli, 1795)—A new sclerochronological archive for the
- 23 Mediterranean? *Journal of Sea Research*, 95, 139-148.
- 24 Calderon-Aguilera, L. E., Aragón-Noriega, E. A., Hand, C. M., & Moreno-Rivera, V. M. (2010).
- 25 Morphometric relationships, age, growth, and mortality of the geoduck clam, *Panopea*
- 26 *generosa*, along the Pacific coast of Baja California, Mexico. *Journal of Shellfish Research*,
- 27 29(2), 319-326.
- 28 Campbell, A., Bourne, N., & Carolsfeld, W. (1990). Growth and size at maturity of the Pacific gaper
- 29 *Tresus nuttallii* (Conrad 1837) in southern British Columbia. *Journal of Shellfish Research*,
- 30 9(2), 273-278.
- 31 Campbell, A., & Ming, M. (2003). Maturity and growth of the Pacific geoduck clam, *Panopea abrupta*,
- 32 in southern British Columbia, Canada. *Journal of Shellfish Research*, 22(1), 85-90.
- 33 Campbell, A., Yeung, C., Dovey, G., & Zhang, Z. (2004). Population biology of the Pacific geoduck
- 34 clam, *Panopea abrupta*, in experimental plots, southern British Columbia, Canada. *Journal of*
- 35 *Shellfish Research*, 23(3), 661-674.
- 36 Campos, E. R., Pena, J. C., Cruz, R. A., & Palacios, J. A. (1998). Growth and reproductive cycle of
- 37 *Polymesoda radiata* (Bivalvia : Corbiculidae) in Costa Rica. *Revista De Biología Tropical*,
- 38 46(3), 643-648.
- 39 Cardoso, J. F., Witte, J. I., & van der Veer, H. W. (2007). Habitat related growth and reproductive
- 40 investment in estuarine waters, illustrated for the tellinid bivalve *Macoma balthica* (L.) in the
- 41 western Dutch Wadden Sea. *Marine Biology*, 152(6), 1271-1282.
- 42 Cardoso, R., & Veloso, V. (2003). Population dynamics and secondary production of the wedge clam
- 43 *Donax hanleyanus* (Bivalvia: Donacidae) on a high-energy, subtropical beach of Brazil.
- 44 *Marine Biology*, 142(1), 153-162. doi:10.1007/s00227-002-0926-2
- 45 Carroll, M. L., Ambrose, W. G., Levin, B. S., Ryan, S. K., Ratner, A. R., Henkes, G. A., & Greenacre, M. J.
- 46 (2011). Climatic regulation of *Clinocardium ciliatum* (bivalvia) growth in the northwestern
- 47 Barents Sea. *Palaeogeography Palaeoclimatology Palaeoecology*, 302(1-2), 10-20.
- 48 doi:10.1016/j.palaeo.2010.06.001
- 49 Carroll, M. L., Ambrose, W. G., Locke V, W. L., Ryan, S. K., & Johnson, B. J. (2014). Bivalve growth rate
- 50 and isotopic variability across the Barents Sea Polar Front. *Journal of Marine Systems*, 130,
- 51 167-180. doi:10.1016/j.jmarsys.2013.10.006
- 52 Cataldo, D., & Boltovskoy, D. (1998). Population dynamics of *Corbicula fluminea* (Bivalvia) in the
- 53 Paraná river delta (Argentina). *Hydrobiologia*, 380(1-3), 153-163.
- 54
- 55
- 56
- 57
- 58
- 59
- 60



- Cayré, P. (1978). Etude de la moule *Perna perna* L. et des possibilités de mytiliculture en République Populaire du Congo. *Cah ORSTOM serie Oceanographie*, 16(1) 9-17.
- Ceccherelli, V. U., & Rossi, R. (1984). Settlement, growth and production of the mussel *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, 16(1), 173-184.
- Chauvaud, L., Patry, Y., Jolivet, A., Cam, E., Le Goff, C., Strand, Ø., . . . Clavier, J. (2012). Variation in Size and Growth of the Great Scallop *Pecten maximus* along a Latitudinal Gradient. *Plos One*, 7(5), e37717. doi:10.1371/journal.pone.0037717
- Chiantore, M., Cattaneo-Vietti, R., & Heilmayer, O. (2003). Antarctic scallop (*Adamussium colbecki*) annual growth rate at Terra Nova Bay. *Polar Biology*, 26(6), 416-419.
- Choo, P., & Speiser, G. (1979). Estimation of the growth parameters and mortality of *Mytilus viridis* Linnaeus (Mollusca, Mytilidae) cultured in a suspended plastic cage in Jelutong, Penang. *Malaysian agricultural journal*.
- Chrysanthakopoulou, V., & Kaspiris, P. (2005). Age and growth of the carpet shell clam *Ruditapes decussatus* (Linnaeus 1758) in Araxos lagoon (NW Peloponnisos, Greece) (Vol. 14).
- Chung, E.-Y., Ryou, D.-K., & Lee, J.-H. (1994). Gonadal development, age and growth of the shortnecked clam, *Ruditapes philippinarum* (Pelecypoda: Veneridae), on the coast of Kimje, Korea. *The Korean Journal of Malacology*, 10(1), 38-48.
- Chute, A. S., McBride, R. S., Emery, S. J., & Robillard, E. (2016). Annulus formation and growth of Atlantic surfclam (*Spisula solidissima*) along a latitudinal gradient in the western North Atlantic Ocean. *Journal of Shellfish Research*, 35(4), 729-737. doi:10.2983/035.035.0402
- Ciocco, N. F. (1991). Differences in individual growth rate among scallop (*Chlamys tehuatlensis* (d'Orb.)) populations from San José Gulf (Argentina). *Fisheries Research*, 12(1), 31-42. doi:https://doi.org/10.1016/0165-7836(91)90047-J
- Claereboudt, M., & Himmelman, J. (1996). Recruitment, growth and production of giant scallops (*Placopecten magellanicus*) along an environmental gradient in Baie des Chaleurs, eastern Canada. *Marine Biology*, 124(4), 661-670.
- Clasing, E., Brey, T., Stead, R., Navarro, J., & Asencio, G. (1994). Population-Dynamics of *Venus antiqua* (Bivalvia, Veneracea) in the Bahia-De-Yaldad, Isla-De-Chiloe, Southern Chile. *Journal of Experimental Marine Biology and Ecology*, 177(2), 171-186.
- Çolakoglu, S. (2014). Population Structure, Growth and Production of the Wedge Clam *Donax trunculus* (Bivalvia, Donacidae) in the West Marmara Sea, Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 14(1), 221-230. doi:10.4194/1303-2712-v14\_1\_24
- Çolakoğlu, S., & Palaz, M. (2014). Some population parameters of *Ruditapes philippinarum* (Bivalvia, Veneridae) on the southern coast of the Marmara Sea, Turkey. *Helgoland Marine Research*, 68(4), 539.
- Cole, H. (1956). A preliminary study of growth-rate in cockles (*Cardium edule* L.) in relation to commercial exploitation. *ICES Journal of Marine Science*, 22(1), 77-90.
- Conan, G., & Shafee, M. S. (1978). Growth and biannual recruitment of the black scallop *Chlamys varia* (L.) in Lanveoc area, Bay of Brest. *Journal of Experimental Marine Biology and Ecology*, 35(1), 59-71.
- Corte, G. N., Coleman, R. A., & Amaral, A. C. Z. (2017). Environmental influence on population dynamics of the bivalve *Anomalocardia brasiliensis*. *Estuarine, Coastal and Shelf Science*, 187, 241-248. doi:https://doi.org/10.1016/j.ecss.2017.01.016
- Cranfield, H., Michael, K., & Francis, R. (1996). Growth rates of five species of subtidal clam on a beach in the South Island, New Zealand. *Marine and Freshwater Research*, 47(6), 773-784. doi:https://doi.org/10.1071/MF9960773
- Cranfield, H. J., & Michael, K. P. (2001). Growth rates of five species of surf clams on a southern North Island beach, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35(5), 909-924.

- Cranford, P., Peer, D., & Gordon, D. (1985). Population dynamics and production of *Macoma balthica* in Cumberland Basin and Shepody Bay, Bay of Fundy. *Netherlands Journal of Sea Research*, 19(2), 135-146.
- Cruz-Vásquez, R., Rodríguez-Domínguez, G., Alcántara-Razo, E., & Aragón-Noriega, E. A. (2012). Estimation of individual growth parameters of the Cortes geoduck *Panopea globosa* from the central Gulf of California using a multimodel approach. *Journal of Shellfish Research*, 31(3), 725-732.
- da Silva, C. F., Corte, G. N., Yokoyama, L. Q., Abrahão, J. R., & Amaral, A. C. Z. (2015). Growth, mortality, and reproduction of *Tagelus plebeius* (Bivalvia: Solecurtidae) in Southeast Brazil. *Helgoland Marine Research*, 69(1), 1-12. doi:10.1007/s10152-014-0417-0
- Dalgıç, G., Okumuş, İ., & Karayücel, S. (2010). The effect of fishing on growth of the clam *Chamelea gallina* (Bivalvia: Veneridae) from the Turkish Black Sea coast. *Journal of the Marine Biological Association of the United Kingdom*, 90(2), 261-265.
- Dang, C., de Montaudouin, X., Gam, M., Paroissin, C., Bru, N., & Caill-Milly, N. (2010). The Manila clam population in Arcachon Bay (SW France): Can it be kept sustainable? *Journal of Sea Research*, 63(2), 108-118. doi:10.1016/j.seares.2009.11.003
- Dare, P. (1991). *Use of external shell microgrowth patterns for determining growth and age in the scallop, Pecten maximus*. Paper presented at the Proc. Eighth International Pectinid Workshop, Cherbourg.
- Dare, P., & Deith, M. (1990). Age determination of scallops, *Pecten maximus* (Linnaeus, 1758), using stable oxygen isotope analysis, with some implications for fisheries management in British waters. *An International Compendium of Scallop Biology and Culture*. World Aquaculture Society, Baton Rouge, 118-133.
- Davenport, J., Davenport, J., & Davies, G. (1984). A preliminary assessment of growth rates of mussels from the Falkland Islands (*Mytilus chilensis* Hupé and *Aulacomya ater* (Molina)). *Ices Journal of Marine Science*, 41(2), 154-158. doi:10.1093/icesjms/41.2.154
- Defeo, O. (1996). Experimental management of an exploited sandy beach bivalve population. *Revista Chilena De Historia Natural*, 69(4), 605-614.
- Defeo, O., Eduardo, J., & Lyonet, A. (1992). Community Structure and Intertidal Zonation of the Macroinfauna on the Atlantic Coast of Uruguay. *Journal of Coastal Research*, 8(4), 830-839.
- Defeo, O., & Gutierrez, N. (2003). Geographical patterns in growth estimates of the scallop *Zygochlamys patagonica*, with emphasis on Uruguayan waters. *Journal of Shellfish Research*, 22(3), 643-646.
- Defeo, O., Ortiz, E., & Castilla, J. C. (1992). Growth, mortality and recruitment of the yellow clam *Mesodesma mactroides* on Uruguayan beaches. *Marine Biology*, 114(3), 429-437. doi:10.1007/BF00350034
- Degraer, S., Meire, P., & Vincx, M. (2007). Spatial distribution, population dynamics and productivity of *Spisula subtruncata*: implications for *Spisula* fisheries in seaduck wintering areas. *Marine Biology*, 152(4), 863-875. doi:10.1007/s00227-007-0740-y
- Dekker, R., & Beukema, J. (1999). Relations of summer and winter temperatures with dynamics and growth of two bivalves, *Tellina tenuis* and *Abra tenuis*, on the northern edge of their intertidal distribution. *Journal of Sea Research*, 42(3), 207-220.
- del Norte, A. C., & Villarta, K. A. (2010). Use of Population Parameters in Examining Changes in the Status of the Short-Necked Clam *Paphia undulata* Born, 1778 (Mollusca, Pelecypoda; Veneridae) in Coastal Waters of Southern Negros Occidental. *Science Diliman*, 22(1), 53-60.
- Del Norte, A. G. C. (1988). Aspects of the growth, recruitment, mortality and reproduction of the scallop *Amusium pleuronectes* (Linné) in the Lingayen Gulf, Philippines. *Ophelia*, 29(2), 153-168. doi:10.1080/00785326.1988.10430826
- Deval, M., & Oray, I. (1998). The annual shell increments of Bivalvia *Chamelea gallina* L. 1758 in the northern Sea of Marmara. *Oebalia*, 24, 93-109.

- Deval, M. C., & Göktürk, D. (2008). Population structure and dynamics of the cut trough shell *Spisula subtruncata* (da Costa) in the Sea of Marmara, Turkey. *Fisheries Research*, 89(3), 241-247. doi:https://doi.org/10.1016/j.fishres.2007.09.019
- Deval, M. C. (2009). Growth and reproduction of the wedge clam (*Donax trunculus*) in the Sea of Marmara, Turkey. *Journal of Applied Ichthyology*, 25(5), 551-558. doi:doi:10.1111/j.1439-0426.2009.01258.x
- Devillers, N. (1998). A comparison of four growth models for evaluating growth of the northern quahog *Mercenaria mercenaria* (L.). *Journal of Shellfish Research*, 17, 191-194.
- Dolmer, P. (1998). The interactions between bed structure of *Mytilus edulis* L. and the predator *Asterias rubens* L. *Journal of Experimental Marine Biology and Ecology*, 228(1), 137-150.
- Edward, J. P., & Ayyakkannu, K. (1991). Temporal variation in annual production of *Tellina nobilis* and *Tellina cuspis* in a tropical estuarine environment. *Mahasagar*, 24(1), 21-29.
- Escati-Peñaloza, G., Parma, A. M., & Orensanz, J. M. (2010). Analysis of longitudinal growth increment data using mixed-effects models: Individual and spatial variability in a clam. *Fisheries Research*, 105(2), 91-101. doi:https://doi.org/10.1016/j.fishres.2010.03.007
- Etim, L., Sankare, Y., Brey, T., & Arntz, W. (1998). The dynamics of unexploited population of *Corbula trigona* (Bivalvia : Corbulidae) in a brackish-water lagoon, Cote d'Ivoire. *Archive of Fishery and Marine Research*, 46(3), 253-262.
- Eversole, A. G., Devillers, N., & Anderson, W. D. (2000). Age and size of *Mercenaria mercenaria* in two sisters creek, South Carolina. *Journal of Shellfish Research*, 19(1), 51-56.
- Ezgeta-Balic, D., Peharda, M., Richardson, C. A., Kuzmanic, M., Vrgoc, N., & Isajlovic, I. (2011). Age, growth, and population structure of the smooth clam *Callista chione* in the eastern Adriatic Sea. *Helgoland Marine Research*, 65(4), 457-465. doi:10.1007/s10152-010-0235-y
- Fahy, E., Carroll, J., O'Toole, M., & Hickey, J. (2003). A preliminary account of fisheries for the surf clam *Spisula solida* (L)(Mactracea) in Ireland (0332-4338). *Irish Fisheries Bulletin*, Marine Institute.
- Fahy, E., & Gaffney, J. (2001). Growth statistics of an exploited razor clam (*Ensis siliqua*) bed at Gormanstown, Co Meath, Ireland. *Hydrobiologia*, 465(1), 139-151. doi:10.1023/a:1014580522523
- Felix-Pico, E. F., Ramirez-Rodriguez, M., & Holguin-Quinones, O. (2009). Growth and Fisheries of the Black Ark *Anadara tuberculosa*, a Bivalve Mollusk, in Bahia Magdalena, Baja California Sur, Mexico. *North American Journal of Fisheries Management*, 29(1), 231-236. doi:10.1577/m06-050.1
- Filippenko, D., & Naumenko, E. (2014). Patterns of the growth of soft-shell clam *Mya arenaria* L. (Bivalvia) in shallow water estuaries of the southern Baltic Sea. *Ecohydrology & Hydrobiology*, 14. doi:10.1016/j.ecohyd.2014.03.002
- Fiori, S., & Defeo, O. (2006). Biogeographic patterns in life-history traits of the yellow clam, *Mesodesma mactroides*, in Sandy Beaches of South America. *Journal of Coastal Research*, 22(4), 872-893.
- Fiori, S. M., & Morsan, E. M. (2004). Age and individual growth of *Mesodesma mactroides* (Bivalvia) in the southernmost range of its distribution. *Ices Journal of Marine Science*, 61(8), 1253-1259. doi:10.1016/j.icesjms.2004.07.025
- Flores, L., Licandeo, R., Cubillos, L. A., & Mora, E. (2014). Intra-specific variability in life-history traits of *Anadara tuberculosa* (Mollusca: Bivalvia) in the mangrove ecosystem of the Southern coast of Ecuador. *Revista De Biología Tropical*, 62(2), 473-482. doi:10.15517/rbt.v62i2.8501
- Flores, L. A. (2011). Growth estimation of mangrove cockle *Anadara tuberculosa* (Mollusca: Bivalvia): application and evaluation of length-based methods. *Revista De Biología Tropical*, 59, 159-170.
- Gagayev, S. (1989). Growth and production of mass species of bivalves in Chaun Bay (East Siberian Sea). *Oceanology*, 29, 504-507.



- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
- Gage, J. D. (1994). Recruitment ecology and age structure of deep-sea invertebrate populations. *Reproduction, larval biology, and recruitment of the deep-sea benthos*. Columbia University Press, New York, 223-242.
- Galinou-Mitsoudi, S., Vlahavas, G., & Papoutsis, O. (2006). Population study of the protected bivalve *Pinna nobilis* (Linnaeus, 1758) in Thermaikos Gulf (north Aegean Sea). *Journal of Biological Research*, 5, 47-53.
- Gam, M., de Montaudouin, X., & Bazairi, H. (2010). Population dynamics and secondary production of the cockle *Cerastoderma edule*: A comparison between Merja Zerga (Moroccan Atlantic Coast) and Arcachon Bay (French Atlantic Coast). *Journal of Sea Research*, 63(3-4), 191-201. doi:10.1016/j.seares.2010.01.003
- García-March, J., García-Carrascosa, A., Cantero, A. P., & Wang, Y.-G. (2007). Population structure, mortality and growth of *Pinna nobilis* Linnaeus, 1758 (Mollusca, Bivalvia) at different depths in Moraira bay (Alicante, Western Mediterranean). *Marine Biology*, 150(5), 861-871.
- Garcia-March, J. R., Marquez-Aliaga, A., Wang, Y. G., Surge, D., & Kersting, D. K. (2011). Study of *Pinna nobilis* growth from inner record: How biased are posterior adductor muscle scars estimates? *Journal of Experimental Marine Biology and Ecology*, 407(2), 337-344. doi:10.1016/j.jembe.2011.07.016
- García-March, J. R., Manuel García-Carrascosa, A., & Luis Pena, A. (2002). In situ measurement of *Pinna nobilis* shells for age and growth studies: a new device. *Marine Ecology*, 23(3), 207-217.
- Garcia, F., & Plante, R. (1993). Color marks from bauxite used in a growth estimation method for *Ruditapes decussatus* in the Bay of Fos. *Comptes Rendus De L Academie Des Sciences Serie Iii-Sciences De La Vie-Life Sciences*, 316(2), 121-126.
- Garcia, N., Prieo, A., Alzola, R., & Lodeiros, C. (2003). Growth and size distribution of *Donax denticulatus* (Mollusca : donacidae) in playa brava, Peninsula de Araya, Sucre State, Venezuela. *Revista Cientifica-Facultad De Ciencias Veterinarias*, 13(6), 464-470.
- Gaspar, M., Pereira, A., Vasconcelos, P., & Monteiro, C. C. (2004). Age and growth of *Chamelea gallina* from the Algarve coast (southern Portugal): Influence of seawater temperature and gametogenic cycle on growth rate. *Journal of Molluscan Studies*, 70, 371-377. doi:10.1093/mollus/70.4.371
- Gaspar, M. B., Castro, M., & Monteiro, C. C. (1995). Age and growth rate of the clam, *Spisula solida* L, from a site off Vilamoura, south Portugal, determined from acetate replicas of shell sections. *Scientia Marina*, 59, 87-93.
- Gaspar, M. B., Pereira, A. M., Vasconcelos, P., & Monteiro, C. C. (2004). Age and growth of *Chamelea gallina* from the Algarve coast (Southern Portugal): Influence of seawater temperature and gametogenic cycle on growth rate. *Journal of Molluscan Studies*, 70(4), 371-377. doi:10.1093/mollus/70.4.371
- Gil, G. M., & Thomé, J. W. (2000). Estudo do crescimento em comprimento de *Donax hanleyanus* Philippi, 1847 (Mollusca, Bivalvia, Donacidae). *Biociências*, 8(2), 163-175.
- Golikov, A. N., & Scarlato, O. A. (1970). Abundance, dynamics and production properties of populations of edible bivalves *Mizuhopecten yessoensis* and *Spisula sachalinensis* related to the problem of organization of controllable submarine farms at the Western shores of the Sea of Japan. *Helgoländer wissenschaftliche Meeresuntersuchungen*, 20(1), 498-513. doi:10.1007/bf01609925
- Goong, S. A., & Chew, K. K. (2001). Growth of butter clams, *Saxidomus giganteus* Deshayes, on selected beaches in the State of Washington. *Journal of Shellfish Research*, 20(1), 143-148.
- Goshima, S., Ide, N., Fujiyoshi, Y., Noda, T., & Nakao, S. (1996). Reproductive Cycle and Shell Growth of Transplanted Manila Clam *Ruditapes philippinarum* in Saroma Lagoon. *Nippon Suisan Gakkaishi (Japanese Edition)*, 62, 195-200. doi:10.2331/suisan.62.195
- Gray, A., Seed, R., & Richardson, C. (1997). Reproduction and growth of *Mytilus edulis chilensis* from the Falkland Islands. *Scientia Marina*, 61, 39-48.

- Green, A., & Craig, P. (1999). Population size and structure of giant clams at Rose Atoll, an important refuge in the Samoan Archipelago. *Coral Reefs*, 18(3), 205-211. doi:10.1007/s003380050183
- Green, J. (1957). The growth of *Scrobicularia plana* (da Costa) in the Gwendraeth estuary. *Journal of the Marine Biological Association of the United Kingdom*, 36(1), 41-47.
- Gribben, P. E., & Creese, R. G. (2003). Protandry in the New Zealand geoduck, *Panopea zelandica* (Mollusca, Bivalvia). *Invertebrate Reproduction & Development*, 44(2-3), 119-129. doi:10.1080/07924259.2003.9652562
- Gribben, P. E., & Creese, R. G. (2005). Age, growth, and mortality of the New Zealand geoduck clam, *Panopea zelandica* (Bivalvia : Hiatellidae) in two north island populations. *Bulletin of Marine Science*, 77(1), 119-135.
- Griffiths, R. J. (1981). Population dynamics and growth of the bivalve *Choromytilus meridionalis* (Kr.) at different tidal levels. *Estuarine, Coastal and Shelf Science*, 12(1), 101-118.
- Guerreiro, J. (1998). Growth and production of the bivalve *Scrobicularia plana* in two southern European estuaries. *Vie et Milieu*, 48(2), 121-131.
- Gusev, A., & Rudinskaya, L. (2014). Shell form, growth, and production of *Astarte borealis* (Schumacher, 1817)(Astartidae, Bivalvia) in the southeastern Baltic Sea. *Oceanology*, 54(4), 458-464.
- Gusev, A. A., & Jurgens-Markina, E. M. (2012). Growth and production of the bivalve *Macoma balthica* (Linnaeus, 1758) (Cardiida: Tellinidae) in the southeastern part of the Baltic Sea. *Russian Journal of Marine Biology*, 38(1), 56-63. doi:10.1134/s1063074012010063
- Gwyther, D., & Mcshane, P. E. (1988). Growth-Rate and Natural Mortality of the Scallop *Pecten alba* Tate in Port-Phillip-Bay, Australia, and Evidence for Changes in Growth-Rate after a 20-Year Period. *Fisheries Research*, 6(4), 347-361.
- Hafsaoui, I., Draredja, B., Lasota, R., Como, S., & Magni, P. (2016). Population dynamics and secondary production of *Donax trunculus* (Mollusca, Bivalvia) in the Gulf of Annaba (Northeast Algeria). *Mediterranean Marine Science*, 17(3), 738-750. doi:10.12681/mms.1760
- Hancock, D. A. (1965). Determination of natural mortality and its causes in an exploited population of cockles (*Cardium edule* L.).
- Harding, J. M. (2007). Northern quahog (= hard clam) *Mercenaria mercenaria* age at length relationships and growth patterns in the York River, Virginia 1954 to 1970. *Journal of Shellfish Research*, 26(1), 101-107.
- Harding, J. M., & Mann, R. (2006). Age and growth of wild Suminoe (*Crassostrea ariakensis*, Fugita 1913) and Pacific (*C. gigas*, Thunberg 1793) oysters from Laizhou Bay, China. *Journal of Shellfish Research*, 25(1), 73-82. doi:10.2983/0730-8000(2006)25[73:aagows]2.0.co;2
- Hart, A. M., & Joll, L. M. (2006). Growth, mortality, recruitment and sex-ratio in wild stocks of silver-lipped pearl oyster *Pinctada maxima* (Jameson)(Mollusca: Pteriidae), in western Australia. *Journal of Shellfish Research*, 25(1), 201-211.
- Haynes, E. B., & Hitz, C. R. (1971). Age and growth of the giant Pacific sea scallop, *Patinopecton caurinus*, from the Strait of Georgia and outer Washington coast. *Journal of the Fisheries Board of Canada*, 28(9), 1335-1341.
- Heald, D., & Caputi, N. (1981). Some aspects of growth, recruitment and reproduction in the southern saucer scallop, *Amusium balloti* (Bernardi, 1861) in Shark Bay, Western Australia. *Fisheries Research Bulletin Western Australia (Australia)*. no. 25.
- Heilmayer, O. (2003). *Environment, adaptation and evolution: scallop ecology across the latitudinal gradient*. (PhD), Universitat Bremen, Bremen, Germany.
- Heilmayer, O., Brey, T., Chiantore, M., Cattaneo-Vietti, R., & Arntz, W. E. (2003). Age and productivity of the Antarctic scallop, *Adamussium colbecki*, in Terra Nova Bay (Ross Sea, Antarctica). *Journal of Experimental Marine Biology and Ecology*, 288(2), 239-256. doi:10.1016/s0022-0981(03)00020-0

- 1
- 2
- 3
- 4 Heilmayer, O., Brey, T., Storch, D., Mackensen, A., & Arntz, W. E. (2004). Population dynamics and
- 5 metabolism of *Aequipecten opercularis* (L.) from the western English Channel (Roscoff,
- 6 France). *Journal of Sea Research*, 52(1), 33-44. doi:DOI 10.1016/j.sears.2003.07.005
- 7 Henderson, S., & Richardson, C. (1994). A comparison of the age, growth rate and burrowing
- 8 behaviour of the razor clams, *Ensis siliqua* and *E. ensis*. *Journal of the Marine Biological*
- 9 *Association of the United Kingdom*, 74(4), 939-954.
- 10 Hendriks, I. E., Basso, L., Deudero, S., Cabanellas-Reboredo, M., & Álvarez, E. (2012). Relative growth
- 11 rates of the noble pen shell *Pinna nobilis* throughout ontogeny around the Balearic Islands
- 12 (Western Mediterranean, Spain). *Journal of Shellfish Research*, 31(3), 749-756.
- 13 Henriques, M. B., & Casarini, L. M. (2009). Growth evaluation of brown mussel *Perna perna* and
- 14 invasive bivalve *Isochnomon bicolor* of a natural bed in Palmas Island, Santos Bay, São Paulo
- 15 State, Brazil. *Boletim Do Instituto De Pesca*, 35(4), 577-586.
- 16 Hermann, M., Carstensen, D., Fischer, S., Laudien, J., Penchaszadeh, P., & Arntz, W. (2009).
- 17 Population structure, growth, and production of the wedge clam *Donax hanleyanus*
- 18 (bivalvia: Donacidae) from Argentinean beaches. *Journal of Shellfish Research*, 28(3), 511-526.
- 19 Hernández-Otero, A., Gaspar, M. B., Macho, G., & Vázquez, E. (2014). Age and growth of the sword
- 20 razor clam *Ensis arcuatus* in the Ría de Pontevedra (NW Spain): Influence of environmental
- 21 parameters. *Journal of Sea Research*, 85, 59-72.
- 22 doi:https://doi.org/10.1016/j.seares.2013.09.006
- 23 Hibbert, C. (1977). Growth and survivorship in a tidal-flat population of the bivalve *Mercenaria*
- 24 *mercenaria* from Southampton water. *Marine Biology*, 44(1), 71-76.
- 25 Hidalgo-De-La-Toba, J. A., Gonzalez-Pelaez, S. S., Morales-Bojórquez, E., Bautista-Romero, J. J., &
- 26 Lluch-Cota, D. B. (2015). Geoduck *Panopea generosa* growth at its southern distribution limit
- 27 in North America using a multimodel inference approach. *Journal of Shellfish Research*,
- 28 34(1), 91-99.
- 29 Higgs, N. D., Reed, A. J., Hooke, R., Honey, D. J., Heilmayer, O., & Thatje, S. (2009). Growth and
- 30 reproduction in the Antarctic brooding bivalve *Adacnarca nitens* (Philobryidae) from the
- 31 Ross Sea. *Marine Biology*, 156(5), 1073-1081. doi:10.1007/s00227-009-1154-9
- 32 Hily, C., & Le Bris, H. (1984). Dynamics of an *Abra alba* population (Bivalve-Scrobiculariidae) in the
- 33 Bay of Brest. *Estuarine, Coastal and Shelf Science*, 19(4), 463-475.
- 34 Hoffmann, A., Bradbury, A., & Goodwin, C. L. (2000). Modeling geoduck, *Panopea abrupta* (Conrad,
- 35 1849) population dynamics. I. Growth. *Journal of Shellfish Research*, 19, 57-62.
- 36 Hopkins, H. S. (1930). Age differences and the respiration in muscle tissues of mollusks. *Journal of*
- 37 *Experimental Zoology*, 56(2), 209-239.
- 38 Hughes, S. E., & Bourne, N. (1981). Stock assessment and life history of a newly discovered Alaska
- 39 surf clam (*Spisula polynyma*) resource in the southeastern Bering Sea. *Canadian Journal of*
- 40 *Fisheries and Aquatic Sciences*, 38(10), 1173-1181.
- 41 Humphreys, J., Caldow, R. W. G., McGrorty, S., West, A. D., & Jensen, A. C. (2007). Population
- 42 dynamics of naturalised Manila clams *Ruditapes philippinarum* in British coastal waters.
- 43 *Marine Biology*, 151(6), 2255. doi:10.1007/s00227-007-0660-x
- 44 Hutchings, J., & Haedrich, R. (1984). Growth and population structure in two species of bivalves
- 45 (Nuculanidae) from the deep sea. *Marine Ecology Progress Series. Oldendorf*, 17(2), 135-142.
- 46 Ignell, S., & Haynes, E. (2000). Geographic patterns in growth of the giant Pacific sea scallop.
- 47 *Patinopecten caurinus*. Fishery Bulletin – National Oceanic and Atmospheric Administration,
- 48 98(4), 849-853.
- 49 Jagadis, I., & Rajagopal, S. (2007). Age and growth of the venus clam *Gafrarium tumidum* (Roding)
- 50 from south-east coast of India. *Indian Journal of Fisheries*, 54(4), 351-356.
- 51 Jiménez, M., Prieto, A., & Ruiz, L. (2004). Distribución de tallas, crecimiento y mortalidad de *Anadara*
- 52 *notabilis* (Bivalvia: Arcidae) en la Bahía de Mochima, Estado Surce, Venezuela. *Boletín del*
- 53 *Instituto Oceanográfico de Venezuela*, 43(1 & 2).
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- Johannessen, O. H. (1973). Length and weight relationships and the potential production of the bivalve *Venerupis pullastra* (Montagu) on a sheltered beach in western Norway. *Sarsia*, 53(1), 41-48.
- Jones, D. S., Arthur, M. A., & Allard, D. J. (1989). Sclerochronological records of temperature and growth from shells of *Mercenaria mercenaria* from Narragansett Bay, Rhode Island. *Marine Biology*, 102(2), 225-234.
- Jones, D. S., Quitmyer, I. R., Arnold, W. S., & Marelli, D. C. (1990). Annual shell banding, age, and growth rate of hard clams (*Mercenaria* spp.) from Florida. *Journal of Shellfish Research*(1).
- Joseph, M. M., & Joseph, P. S. (1985). Age and growth of the oyster *Crassostrea madrasensis* (Preston) in Mulki estuary, west coast of India.
- Juric, I., Buselic, I., Ezgeta-Balic, D., Vrgoc, N., & Peharda, M. (2012). Age, Growth and Condition Index of *Venerupis decussata* (Linnaeus, 1758) in the Eastern Adriatic Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 12(3), 613-618. doi:10.4194/1303-2712-v12\_3\_08
- Kafanov, A. (1985). Growth and production of the bivalve mollusc *Macoma balthica* in the Nabil Lagoon (The Northeastern Sakhalin). *Biologiya Morya Marine Biology*(6), 23-31.
- Kandeel, K. (2016). Population analysis of *Tapes decussata* (Bivalvia: Veneridae) in Lake Timsah, Suez Canal, Egypt. *Journal of Advances in Biology*, 9(1).
- Kandeel, K. E., Mohammed, S. Z., Mostafa, A. M., & Abd-Alla, M. E. (2017). Population Dynamics of the Cockle *Cerastoderma glaucum*: A comparison between Lake Qarun and Lake Timsah, Egypt. *Turkish Journal of Fisheries and Aquatic Sciences*, 17(5), 945-958. doi:10.4194/1303-2712-v17\_5\_10
- KANG, Y. J., & KIM, C. K. (1983). Studies on the structure and production processes of biotic communities in the coastal shallow waters of Korea 3. Age and growth of *Spisula sachalinensis* from the eastern waters of Korea. *Korean Journal of Fisheries and Aquatic Sciences*, 16(2), 82-87.
- KATO, Y., & HAMAI, I. (1975). Growth and shell formation of the surf clam, *Spisula sachalinensis* (Schrenck). *Bulletin of the Faculty of Fisheries, Hokkaido University*, 25(4), 291-303.
- Kennish, M., & Loveland, R. (1980). *Growth models of the northern quahog, Mercenaria mercenaria* (Linné). Paper presented at the Proceedings of the National Shellfisheries Association.
- Kevrekidis, T., & Koukouras, A. (1992). Population dynamics, growth and productivity of *Abra ovata* (Mollusca, Bivalvia) in the Evros delta (North Aegean Sea). *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 77(2), 291-301.
- Khan, M. A., Assim, Z. B., & Ismail, N. (2010). Population dynamics of the green-lipped mussel, *Perna viridis* from the Offshore waters of Naf River coast, Bangladesh. *Chiang Mai Journal of Science*, 37, 344-354.
- Kilada, R. W., Campana, S. E., & Roddick, D. (2006). Validated age, growth, and mortality estimates of the ocean quahog (*Arctica islandica*) in the western Atlantic. *ICES Journal of Marine Science*, 64(1), 31-38.
- Kilada, R. W., Campana, S. E., & Roddick, D. (2009). Growth and sexual maturity of the northern propellerclam (*Cyrtodaria siliqua*) in Eastern Canada, with bomb radiocarbon age validation. *Marine Biology*, 156(5), 1029-1037. doi:10.1007/s00227-009-1146-9
- Kilada, R. W., Roddick, D., & Mombourquette, K. (2007). Age determination, validation, growth and minimum size of sexual maturity of the Greenland smoothcockle (*Serripes groenlandicus*, Bruguiere, 1789) in eastern Canada. *Journal of Shellfish Research*, 26(2), 443-451.
- Kim, B., Ko, T., Song, H., Lee, S., & Kim, S. (1985). Studies on the spawning, and growth of hen clam, *Macra sulcataria* (REEVE). *Bulletin of Fisheries Research & Development Agency*, 34, 157-164.
- Kim, J.-H., Kim, J.-S., Kim, Y.-H., Chung, E.-Y., & Ryu, D.-K. (2003). Age and growth of the jedo venus clam, *Protothaca jedoensis* on the west coast of Korea. *The Korean Journal of Malacology*, 19(2), 125-132.



- Kitamura, A. (2011). Age and growth of *Glossocardia obesa*, a large bivalve in a submarine cave within a coral reef, as revealed by oxygen isotope analysis. *Veliger*, 51, 59-65.
- Kithsiri, H. M., Wijeyaratne, M. J. S., & Amarasinghe, U. (2004). Population dynamics of three commercially important bivalve species (Family: Veneridae) in Puttalam lagoon and Dutch bay, Sri Lanka. *Sri Lanka Journal of Aquatic Sciences*, 9, 13-30. doi:10.4038/sljas.v9i1.7463
- Kraus, M., Beal, B., Chapman, S., & McMartin, L. (1992). A comparison of growth rates in *Arctica islandica* (Linnaeus, 1767) between field and laboratory populations. *Journal of Shellfish Research*, 11, 289-289.
- Kunitzer, A. (1989). Factors affecting the population dynamics of *Amphiura filiformis* (Echinodermata: Ophiuroidea) and *Mysella bidentata* (Bivalvia: Galeommatacea) in the North Sea. *Reproduction, Genetics and Distributions of Marine Organisms* (JS Ryland & P. Tyler, eds), 395-406.
- Lambert, J., & Préfontaine, G. (1995). *The Iceland Scallop, Chlamys Islandica*, in Nunavik: Department of Fisheries & Oceans, Invertebrates & Experimental Biology Division, Maurice Lamontagne Institute.
- Landry, T., Sephton, T., & Jones, D. A. (1993). Growth and mortality of northern quahog, (Linnaeus, 1758) *Mercenaria mercenaria* in Prince Edward Island. *Journal of Shellfish Research*, 12(2), 321-327.
- Langton, R. W., Robinson, W. E., & Schick, D. (1987). Fecundity and reproductive effort of sea scallops *Placopecten magellanicus* from the Gulf of Maine. *Marine Ecology Progress Series*, 19-25.
- Lasta, M. (2001). Argentinian shelf. Part II: Population dynamics of *Zygochlamys patagonica*. *Archive of Fishery and Marine Research*, 49, 125-137..
- Lastra, M., Palacio, J., & Mora, J. (1993). Population dynamics and secondary production of *Abra alba* (Wood) (Bivalvia) in the Santander Bay, northern Spain. *Sarsia*, 78(1), 35-42. doi:10.1080/00364827.1993.10413520
- Laudien, J., Brey, T., & Arntz, W. E. (2003). Population structure, growth and production of the surf clam *Donax serra* (Bivalvia, Donacidae) on two Namibian sandy beaches. *Estuarine Coastal and Shelf Science*, 58, 105-115. doi:10.1016/s0272-7714(03)00044-1
- Laxmilatha, P. (2013). *Population dynamics of the edible clam Meretrix casta* (Chemnitz) (International Union for Conservation of Nature status: Vulnerable) from two estuaries of North Kerala, south west coast of India (Vol. 5).
- Lee, A. M., Williams, A. J., & Southgate, P. C. (2008). Modelling and comparison of growth of the silver-lip pearl oyster *Pinctada maxima* (Jameson) (Mollusca : Pteriidae) cultured in West Papua, Indonesia. *Marine and Freshwater Research*, 59(1), 22-31. doi:https://doi.org/10.1071/MF07112
- Lee, S. Y. (1985). The population dynamics of the green mussel, *Perna viridis* (L. ) in Victoria Harbour, Hong Kong—dominance a polluted environment. *Asian Marine Biology*, 2, 107-118.
- Lee, T. (1973). Scallop and queen scallop survey along the east and south-east coasts of Ireland—August 1972. *Fisheries Development Division, Bord Lascaigh Mhara* 23.
- Lefort, Y. (1994). Growth and mortality of the tropical scallops: *Annachlamys flabellata* (Bernardi), *Comptopallium radula* (Linne) and *Mimachlamys gloriosa* (Reeve) in Southwest Lagoon of New Caledonia. *Journal of Shellfish Research*, 13(2), 539-546.
- Leontarakis, P., & Richardson, C. (2005). Growth of the smooth clam, *Callista chione* (Linnaeus, 1758)(Bivalvia: Veneridae) from the Thracian Sea, northeastern Mediterranean. *Journal of Molluscan Studies*, 71(2), 189-192.
- Lewis, C. V. W., Weinberg, J. R., & Davis, C. S. (2001). Population structure and recruitment of the bivalve *Arctica islandica* (Linnaeus, 1767) on Georges Bank from 1980-1999. *Journal of Shellfish Research*, 20(3), 1135-1144.

- 1
- 2
- 3
- 4 Lim, H.-S., & Lee, C.-I. (2004). Growth Pattern of Soft Clam (*Mya arenaria oonogai*) (Mollusca:
- 5 Bivalvia) from a Mud Flat on the Southwest Coast of Korea. *Korean Journal of Fisheries and*
- 6 *Aquatic Sciences*, 37(2) 105-115.
- 7 Lizarralde, Z., Pittaluga, S., Albarracin, T., & Perroni, M. (2018). Population dynamics and secondary
- 8 production of *Darina solenoides* (Bivalvia: Mactridae) in the Rio Gallegos Estuary, southern
- 9 Patagonia. *Latin American Journal of Aquatic Research*, 46(2), 411-415. doi:10.3856/vol46-
- 10 issue2-fulltext-16
- 11 Lizarralde, Z. I., & Cazzaniga, N. J. (2009). Population dynamics and production of *Tellina petitana*
- 12 (Bivalvia) on a sandy beach of Patagonia, Argentina. *Thalassas*, 25(1), 45-57.
- 13 Llana, M. E. G. (1988). Growth, mortality and recruitment of the Asian moon scallop (*Amusium*
- 14 *pleuronectes*) in the Visayan Sea, Philippines In S. C. Venema, J. M. Christensen, & D. e. Pauly
- 15 (Eds.).
- 16 Lodeiros, C., Rengel, J. J., & Himmelman, J. H. (1999). Growth of *Pteria colymbus* (Roding, 1798) in
- 17 suspended culture in Golfo de Cariaco, Venezuela. *Journal of Shellfish Research*, 18, 155-158.
- 18 Lomovasky, B. J., Baldoni, A., Ribeiro, P., Alvarez, G., Lasta, M., Campodonico, S., & Iribarne, O.
- 19 (2011). Exploring the causes of differences in growth rate of the Patagonian scallop
- 20 *Zygochlamys patagonica* along its commercial bed distribution in the SW Atlantic. *Journal of*
- 21 *Sea Research*, 66(2), 162-171. doi:10.1016/j.seares.2011.05.010
- 22 Lomovasky, B. J., Brey, T., Klugel, A., & Iribarne, O. (2018). Distribution pattern, density and growth
- 23 of the stout razor clam *Tagelus plebeius* in a South-west Atlantic estuarine system. *Journal of*
- 24 *the Marine Biological Association of the United Kingdom*, 98(3), 485-494.
- 25 doi:10.1017/s0025315416001715
- 26 Lomovasky, B. J., Brey, T., & Morriconi, E. (2005). Population dynamics of the venerid bivalve *Tawera*
- 27 *gayi* (Hupe, 1854) in the Ushuaia Bay, Beagle Channel. *Journal of Applied Ichthyology*, 21(1),
- 28 64-69. doi:10.1111/j.1439-0426.2004.00599.x
- 29 Lomovasky, B. J., Lasta, M., Valinas, M., Bruschetti, M., Ribeiro, P., Campodonico, S., & Iribarne, O.
- 30 (2008). Differences in shell morphology and internal growth pattern of the Patagonian
- 31 scallop *Zygochlamys patagonica* in the four main beds across their SW Atlantic distribution
- 32 range. *Fisheries Research*, 89(3), 266-275. doi:10.1016/j.fishres.2007.09.006
- 33 Lomovasky, B. J., Morriconi, E., Brey, T., & Calvo, J. (2002). Individual age and connective tissue
- 34 lipofuscin in the hard clam *Eurhomalea exalbida*. *Journal of Experimental Marine Biology and*
- 35 *Ecology*, 276(1-2), 83-94. doi:10.1016/s0022-0981(02)00240-x
- 36 López-Jamar, E., González, G., & Mejuto, J. (1987). Ecology, growth and production of *Thyasira*
- 37 *flexuosa* (Bivalvia, Lucinacea) from Ría de la Coruña, North-West Spain. *Ophelia*, 27(2), 111-
- 38 126. doi:10.1080/00785236.1987.10422015
- 39 Lucero, C., Cantera, J., & Neira, R. (2012). The fisheries and growth of Ark Clams (Arcoida: Arcidae)
- 40 *Anadara tuberculosa* in Malaga Bay, Colombian Pacific, 2005-2007. *Revista De Biologia*
- 41 *Tropical*, 60(1), 203-217.
- 42 Luckens, P. A. (1990). Distribution, size-frequency, and growth-ring analyses of *Tawera mawsoni*
- 43 (Bivalvia: Veneridae) at Macquarie Island. *New Zealand Journal of Marine and Freshwater*
- 44 *Research*, 24(1), 59-73. doi:10.1080/00288330.1990.9516402
- 45 Luzzatto, D. C. (2007). *Dinámica poblacional de la almeja amarilla (Mesodesma mactroides) y del*
- 46 *berberecho (Donax hanleyanus) en intermareales de playas arenosas del Nordeste de la*
- 47 *Provincia de Buenos Aires*. (Ph.D Thesis), Universidad de Buenos Aires,
- 48 MacDonald, B. A., & Bourne, N. F. (1987). Growth, Reproductive Output, and Energy Partitioning in
- 49 Weathervane Scallops, *Patinopecten caurinus*, from British Columbia. *Canadian Journal of*
- 50 *Fisheries and Aquatic Sciences*, 44(1), 152-160. doi:10.1139/f87-020
- 51 MacDonald, B. A., & Thompson, R. J. (1986). Production, dynamics and energy partitioning in two
- 52 populations of the giant scallop *Placopecten magellanicus* (Gmelin). *Journal of Experimental*
- 53 *Marine Biology and Ecology*, 101(3), 285-299. doi:https://doi.org/10.1016/0022-
- 54 0981(86)90269-8
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3 MacDonald, B. A., Thompson, R. J., & Bourne, N. F. (1991). Growth and Reproductive Energetics of
- 4 three Scallop Species from British Columbia (*Chlamys hastata*, *Chlamys rubida*, and
- 5 *Crassadoma gigantea*). *Canadian Journal of Fisheries and Aquatic Sciences*, 48(2), 215-221.
- 6 doi:10.1139/f91-029
- 7
- 8 Mancera, E., & Mendo, J. (1996). Population dynamics of the oyster *Crassostrea rhizophorae* from
- 9 the Ciénaga Grande de Santa Marta, Colombia. *Fisheries Research*, 26(1), 139-148.
- 10 doi:https://doi.org/10.1016/0165-7836(95)00401-7
- 11 Marcano, J. S., Priet, A., Larez, A., Alio, J. J., & Sanabria, H. (2005). Growth and mortality of *Pinctada*
- 12 *imbricata* (Mollusca : Pteridae) in Guamachito, Araya Peninsula, Sucre State, Venezuela.
- 13 *Ciencias Marinas*, 31(2), 387-397. doi:10.7773/cm.v31i2.55
- 14 Mason, J. (1983). Scallop and queen fisheries in the British Isles. In H. Caspers (Ed.), (Vol. 70, pp.
- 15 440): Farnham (UK) Fishing News Books.
- 16 Mattos, G., & Cardoso, R. S. (2012). Population dynamics of two suspension-feeding bivalves on a
- 17 sheltered beach in southeastern Brazil. *Helgoland Marine Research*, 66(3), 393-400.
- 18 doi:10.1007/s10152-011-0280-1
- 19 Mavuti, K. M., Kimani, E. N., & Mukiana, T. (2005). Growth patterns of the pearl oyster *Pinctada*
- 20 *margaritifera* L. in Gazi Bay, Kenya. *African Journal of Marine Science*, 27(3), 567-575.
- 21 doi:10.2989/18142320509504117
- 22
- 23 Maximovich, N. V., & Guerassimova, A. V. (2003). Life history characteristics of the clam *Mya*
- 24 *arenaria* in the White Sea. *Helgoland Marine Research*, 57(2), 91-99. doi:10.1007/s10152-
- 25 003-0137-3
- 26
- 27 McQuaid, C. D., & Lindsay, T. L. (2000). Effect of wave exposure on growth and mortality rates of the
- 28 mussel *Perna perna*: bottom up regulation of intertidal populations. *Marine Ecology*
- 29 *Progress Series*, 206, 147-154.
- 30 Mendo, J., & Jurado, E. (1993). Length-based growth parameter estimates of the Peruvian scallop
- 31 (*Argopecten purpuratus*). *Fisheries Research*, 15(4), 357-367.
- 32 doi:https://doi.org/10.1016/0165-7836(93)90086-M
- 33
- 34 Merk, V. (2015). *Potential of Greenland cockles (Serripes groenlandicus) as high resolution Arctic*
- 35 *climate archive [Potential der Grönlandherzmuschel (Serripes groenlandicus) als*
- 36 *hochauflösendes Klimaarchiv der Arktis]*. Alfred-Wegener-Institut Helmholtz-Zentrum für
- 37 Polarund Meeresforschung.
- 38 Merrill, A., Posgay, J., & Nichy, F. (1966). Annual marks on shell and ligament of sea scallop
- 39 (*Placopecten magellanicus*). *Fisheries Bulletin*, 65, 299-311.
- 40 Metaxatos, A. (2004). Population dynamics of the venerid bivalve *Callista chione* (L.) in a coastal area
- 41 of the eastern Mediterranean. *Journal of Sea Research*, 52(4), 293-305.
- 42 doi:10.1016/j.seares.2004.03.001
- 43
- 44 Milione, M., & Southgate, P. (2012). Growth of the Winged Pearl Oyster, *Pteria penguin*, at Dissimilar
- 45 Sites in Northeastern Australia. *Journal of Shellfish Research*, 31(1), 13-20.
- 46 doi:10.2983/035.031.0102
- 47 Mirzaei, M. R., Yasin, Z., & Shau Hwai, A. T. (2014). Length-weight relationship, growth and mortality
- 48 of *Anadara granosa* in Penang Island, Malaysia: an approach using length-frequency data
- 49 sets. *Journal of the Marine Biological Association of the United Kingdom*, 95(2), 381-390.
- 50 doi:10.1017/S0025315414001337
- 51 Mistri, M., Rossi, R., & Ceccherelli, V. U. (1988). Growth and Production of the Ark Shell *Scapharca*
- 52 *inaequivalvis* (BRUGUIÈRE) in a Lagoon of the Po River Delta. *Marine Ecology*, 9(1), 35-49.
- 53 doi:doi:10.1111/j.1439-0485.1988.tb00197.x
- 54
- 55 Mohammed, S. Z., & Yassien, M. H. (2003). Population parameters of the pearl oyster *Pinctada*
- 56 *radiata* (Leach) in Qatari waters, Arabian Gulf. *Turkish Journal of Zoology*, 27, 339-343.
- 57 Mohite, S., & Mohite, A. (2009). Age and growth of the shortneck clam, *Paphia malabarica*
- 58 (Chemintz) in estuarine areas of Ratnagiri, West Coast of India. *Asian Journal of Animal*
- 59 *Science*, 3(2), 235-240.
- 60



- Monti, D., Frenkiel, L., & Moueza, M. (1991). Demography and growth of *Anomalocardia brasiliana* (Gmelin) (Bivalvia: Veneridae) in a mangrove in Guadeloupe (French West Indies). *Journal of Molluscan Studies*, 57(2), 249-257. doi:10.1093/mollus/57.2.249
- Morsán, E., & Ciocco, N. F. (2004). Age and growth model for the southern geoduck, *Panopea abbreviata*, off Puerto Lobos (Patagonia, Argentina). *Fisheries Research*, 69(3), 343-348. doi:https://doi.org/10.1016/j.fishres.2004.06.012
- Morsan, E., Zaidman, P., Ocampo-Reinaldo, M., & Ciocco, N. (2010). Population structure, distribution and harvesting of southern geoduck, *Panopea abbreviata*, in San Matías Gulf (Patagonia, Argentina). *Scientia Marina*, 74(4), 763-772. doi:10.3989/scimar.2010.74n4763
- Morsan, E. M., & Orensanz, J. M. L. (2004). Age structure and growth in an unusual population of purple clams, *Amiantis purpuratus* (Lamarck, 1818) (Bivalvia; Veneridae), from Argentine Patagonia. *Journal of Shellfish Research*, 23(1), 73-80.
- Moss, D., Surge, D., & Khaitov, V. (2018). Lifespan and growth of *Astarte borealis* (Bivalvia) from Kandalaksha Gulf, White Sea, Russia. *Polar Biology*. doi:10.1007/s00300-018-2290-9
- Moura, P., Garaulet, L. L., Vasconcelos, P., Chainho, P., Costa, J. L., & Gaspar, M. B. (2017). Age and growth of a highly successful invasive species: the Manila clam *Ruditapes philippinarum* (Adams & Reeve, 1850) in the Tagus Estuary (Portugal). *Aquatic Invasions*, 12(2), 133-146. doi:10.3391/ai.2017.12.2.02
- Moura, P., Gaspar, M. B., & Monteiro, C. C. (2009). Age determination and growth rate of a *Callista chione* population from the southwestern coast of Portugal. *Aquatic Biology*, 5(1), 97-106. doi:10.3354/ab00119
- Moura, P., Vasconcelos, P., & Gaspar, M. B. (2013). Age and growth in three populations of *Dosinia exoleta* (Bivalvia: Veneridae) from the Portuguese coast. *Helgoland Marine Research*, 67(4), 639-652. doi:10.1007/s10152-013-0350-7
- Munch-Petersen, S., & Kristensen, P. S. (2001). On the dynamics of the stocks of blue mussels (*Mytilus edulis* L.) in the Danish Wadden Sea. In G. Burnell (Ed.), *Coastal Shellfish — A Sustainable Resource: Proceedings of the Third International Conference on Shellfish Restoration, held in Cork, Ireland, 28 September–2 October 1999* (pp. 31-43). Dordrecht: Springer Netherlands.
- Murawski, S., W. Ropes, J., & Serchuk, F. M. (1982). Growth of the ocean quahog, *Arctica islandica*, in the Middle Atlantic Bight. *Fisheries Bulletin*, 80(1), 21-34.
- Myasnikov, M. G., & Kochnev, Y. P. (1988). Lifespan, growth and sex structure of scallop (*Chlamys albidus*) off the Kuril Islands. *Marine Commercial Invertebrates. Collected Papers. Moscow: VNIRO*, 153 - 166.
- Naidu, K., Sundraj, F. M., Cahill, F. M., & Lewis, D. B. (1982). *Status and Assessment of the Iceland Scallop, Chlamys Islandica, in the Northeastern Gulf of St. Lawrence*. Canadian Atlantic Fisheries Scientific Advisory Committee.
- Narasimham, K. A. (1988). Biology of the blood clam *Anadara granosa* (Linnaeus) in Kakinada Bay. *Journal of the Marine Biological Association of India*, 30(1&2) 137-150.
- Newcombe, C. L. (1935). Growth of *Mya arenaria* L. in the Bay of Fundy Region. *Canadian Journal of Research*, 13, 97-137. doi:10.1139/cjr35d-009
- Niamaimandi, N. (2014). Growth, Mortality and Stock Abundance of Venerid Bivalve, *Paphia cor* from Iranian Coastal Waters of Bushehr, Persian Gulf. *Environmental Studies of Persian Gulf*, 1(1), 51-58.
- Nickerson, R. B. (1975). *A Critical analysis of some razor clam (Siliqua patula, Dixon) populations in Alaska*: State of Alaska Department of Fish and Game.
- Nolan, C. P. (1988). *Calcification and growth rates in Antarctic molluscs. British Antarctic Survey Report AD6*.
- Nolan, C. P., & Clarke, A. (1993). Growth in the bivalve *Yoldia eightsi* at Signy Island, Antarctica, determined from internal shell increments and calcium-45 incorporation. *Marine Biology*, 117(2), 243-250. doi:10.1007/bf00345669

- Nugranad, J. (1988). *Preliminary report on the growth, mortality and recruitment of the Asian moon scallop Amusium pleuronectes in Koh Chang (Koh Kood area), eastern Gulf of Thailand.* (Unpublished Report).
- Nurul Amin, S., Zafar, M., & Halim, A. (2008). Age, growth, mortality and population structure of the oyster, *Crassostrea madrasensis*, in the Moheshkhali Channel (southeastern coast of Bangladesh). *Journal of Applied Ichthyology*, 24(1), 18-25.
- O'Brien, K., & Keegan, B. F. (2005). Determination of Growth Dynamics of the Bivalve *Abra alba* (Wood) in Kinsale Harbour (S Coast of Ireland) Using Modal Progression Analysis. *The Irish Naturalists' Journal*, 28(3), 97-106.
- Ocaña, F. A. (2015). Growth and production of *Donax striatus* (Bivalvia: Donacidae) from Las Balsas beach, Gibara, Cuba. *Revista De Biología Tropical*, 63, 639-646.
- Ocana, F. A., Apin, Y. C., & Cala, Y. R. (2013). Population dynamic of *Donax denticulatus* (Bivalvia: Donacidae) at Carenero Beach, Southeastern Cuba. *Revista De Biología Tropical*, 61(4), 1637-1646.
- Oker, W. (1976). Observations on some population parameters of exploited stocks of *Senilia senilis* (= *Arca senilis*) in Sierra Leone. *Marine Biology*, 38(3), 217-229. doi:10.1007/bf00388935
- Olivier, S. R., & Instituto de Biología, M. (1971). *Estructura de la comunidad, dinámica de la población y biología de la almeja amarilla (Mesodesma mactroides Desh. 1854) en Mar Azul (Pdo. de Gral. Madariaga, Bs. As., Argentina)*. Mar del Plata: Instituto de Biología Marina.
- Orensanz, J. M. (1986). Size, environment and density : the regulation of a scallop stock and its management implications. *North Pacific Workshop on Stock Assessment and Management of Invertebrates*, 1986.
- Pace, S. M., Powell, E. N., Mann, R., Long, M. C., & Klinck, J. M. (2017). Development of an age-frequency distribution for ocean quahogs (*Arctica islandica*) on Georges Bank. *Journal of Shellfish Research*, 36(1), 41-53. doi:10.2983/035.036.0106
- Palacio, J., Cruz, R. A., & Urpi, O. P. (1983). Estructura poblacional y cuantificación de *Donax dentifer* Hanley, 1843 (Pelecypoda: Donacidae) en Playa Garza, Puntarenas, Costa Rica. *Revista De Biología Tropical*, 31(2), 251 - 255.
- Palmer, D. W. (2004). Growth of the razor clam *Ensis directus*, an alien species in the Wash on the east coast of England. *Journal of the Marine Biological Association of the UK*, 84, 1075-1076. doi:10.1017/S0025315404010458h
- Pearson, R., & Munro, J. (1991). Growth, mortality and recruitment rates of giant clams, *Tridacna gigas* and *T. derasa*, at Michaelmas Reef, central Great Barrier Reef, Australia. *Marine and Freshwater Research*, 42(3), 241-262. doi:https://doi.org/10.1071/MF9910241
- Pedersen, S. (1994). Population Parameters of the Iceland Scallop (*Chlamys islandica* (Müller)) from West Greenland. *Journal of Northwest Atlantic Fishery Science*, 16, 75-87. doi:10.2960/J.v16.a7
- Peharda, M., Crnčević, M., Bušelić, I., Richardson, C. A., & Ezgeta-Balić, D. (2012). Growth And Longevity of *Glycymeris nummaria* (Linnaeus, 1758) from the Eastern Adriatic, Croatia. *Journal of Shellfish Research*, 31(4), 947-950, 944.
- Peharda, M., Ezgeta-Balić, D., Radman, M., Sinjkević, N., Vrgoč, N., & Isajlović, I. (2011). Age, growth and population structure of *Acanthocardia tuberculata* (Bivalvia: Cardiidae) in the eastern Adriatic Sea. 2011, 76(1), 8. doi:10.3989/scimar.03257.21A
- Peharda, M., Popovic, Z., Ezgeta-Balic, D., Vrgoc, N., Puljas, S., & Frankic, A. (2013). Age and growth of *Venus verrucosa* (Bivalvia: Veneridae) in the eastern Adriatic Sea. *Cahiers De Biologie Marine*, 54(2), 281-286.
- Peharda, M., Puljas, S., Chauvaud, L., Schone, B. R., Ezgeta-Balic, D., & Thebault, J. (2015). Growth and longevity of *Lithophaga lithophaga*: what can we learn from shell structure and stable isotope composition? *Marine Biology*, 162(8), 1531-1540. doi:10.1007/s00227-015-2690-0

- Peharda, M., Richardson, C. A., Mladineo, I., Šestanović, S., Popović, Z., Bolotin, J., & Vrgoč, N. (2007). Age, growth and population structure of *Modiolus barbatus* from the Adriatic. *Marine Biology*, 151(2), 629-638. doi:10.1007/s00227-006-0501-3
- Peharda, M., Richardson, C. A., Onofri, V., Bratoš, A. N. A., & Crnčević, M. (2002). Age and growth of the bivalve *Arca noae* L. in the Croatia Adriatic Sea. *Journal of Molluscan Studies*, 68(4), 307-310. doi:10.1093/mollus/68.4.307
- Peharda, M., Soldo, A., Pallaoro, A., Matić, S., & Cetinić, P. (2003). Age and growth of the Mediterranean scallop *Pecten jacobaeus* (Linnaeus 1758) in the northern Adriatic Sea. *Journal of Fisheries Research* 22(3), 639.
- Pérez-Valencia, S. A., & Aragon-Noriega, E. (2013). Age and growth of the Cortes Geoduck *Panopea globosa* (Dall, 1898) in the upper Gulf of California. *Indian Journal of Geo-Marine Sciences*, 42(2), 201-205.
- Perez, V., Olivier, F., Tremblay, R., Neumeier, U., Thébault, J., Chauvaud, L., & Meziane, T. (2013). Trophic resources of the bivalve, *Venus verrucosa*, in the Chausey archipelago (Normandy, France) determined by stable isotopes and fatty acids. *Aquatic Living Resources*, 26(3), 229-239. doi:10.1051/alr/2013058
- Philipp, E., Brey, T., Heilmayer, O., Abele, D., & Pörtner, H.-O. (2006). Physiological ageing in a polar and a temperate swimming scallop. *Marine Ecology Progress Series*, 307, 187-198.
- Philipp, E., Brey, T., Portner, H. O., & Abele, D. (2005). Chronological and physiological ageing in a polar and a temperate mud clam. *Mechanisms of Ageing and Development*, 126(5), 598-609. doi:10.1016/j.mad.2004.12.003
- Pinn, E. H., Richardson, C. A., Thompson, R. C., & Hawkins, S. J. (2005). Burrow morphology, biometry, age and growth of piddocks (Mollusca: Bivalvia: Pholadidae) on the south coast of England. *Marine Biology*, 147(4), 943-953. doi:10.1007/s00227-005-1582-0
- Ponurovskii, S. K. (2008). Population structure and growth of the Japanese littleneck clam *Ruditapes philippinarum* in Amursky Bay, Sea of Japan. *Russian Journal of Marine Biology*, 34(5), 329-332. doi:10.1134/s1063074008050106
- Pouvreau, S., & Prasil, V. (2001). Growth of the black-lip pearl oyster, *Pinctada margaritifera*, at nine culture sites of French Polynesia: synthesis of several sampling designs conducted between 1994 and 1999. *Aquatic Living Resources*, 14(3), 155-163. doi:10.1016/s0990-7440(01)01120-2
- Pranovi, F., Marcato, S., & Zanellato, R. (1994). Analisi biometriche e biologia di popolazione del mollusco antartico *Adamussium colbecki* a Baia Terra Nova, Mare di Ross. *Atti Dell'Istituto Veneto Di Scienze Lettere Ed Arti*, 152.
- Puljas, S., Peharda, M., Zupan, I., & Buksa, F. (2015). Maximum recorded life span of *Arca noae* Linnaeus, 1758 in the marine protected area Telascica, Adriatic Sea. *Cahiers De Biologie Marine*, 56(2), 163-168.
- Rabaoui, L., Tlig-Zouari, S., Katsanevakis, S., Belgacem, W., & Hassine, O. K. B. (2011). Differences in absolute and relative growth between two shell forms of *Pinna nobilis* (Mollusca: Bivalvia) along the Tunisian coastline. *Journal of Sea Research*, 66(2), 95-103. doi:https://doi.org/10.1016/j.seares.2011.05.002
- Rabaoui, L., Tlig Zouari, S., Katsanevakis, S., & Ben Hassine, O. K. (2007). Comparison of absolute and relative growth patterns among five *Pinna nobilis* populations along the Tunisian coastline: an information theory approach. *Marine Biology*, 152(3), 537-548. doi:10.1007/s00227-007-0707-z
- Ralph, R., & Maxwell, J. G. H. (1977). Growth of two Antarctic lamellibranchs: *Adamussium colbecki* and *Laternula elliptica*. *Marine Biology*, 42(2), 171-175. doi:10.1007/bf00391569
- Ramon, M. (2003). Population dynamics and secondary production of the cockle *Cerastoderma edule* (L.) in a backbarrier tidal flat of the Wadden Sea. *Scientia Marina*, 67(4), 429-443. doi:10.3989/scimar.2003.67n4429

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
- Ramón, M., Abelló, P., & Richardson, C. A. (1995). Population structure and growth of *Donax trunculus* (Bivalvia: Donacidae) in the western Mediterranean. *Marine Biology*, 121(4), 665-671. doi:10.1007/bf00349302
- Ramón, M., & Richardson, C. (1992). Age determination and shell growth of *Chamelea gallina* (Bivalvia: Veneridae) in the western Mediterranean. *Marine Ecology Progress Series*, 15-23.
- Rapson, A. (1952). The Toheroa, *Amphidesma ventricosum* Gray (Eulamellibranchiata), Development and Growth. *Marine and Freshwater Research*, 3(2), 170-198. doi:https://doi.org/10.1071/MF9520170
- Reed, A. J., Linse, K., & Thatje, S. (2014). Differential adaptations between cold-stenothermal environments in the bivalve *Lissarca cf. miliaris* (Philobryidae) from the Scotia Sea islands and Antarctic Peninsula. *Journal of Sea Research*, 88, 11-20.
- Reed, A. J., Morris, J. P., Linse, K., & Thatje, S. (2014). Reproductive morphology of the deep-sea protobranch bivalves *Yoldiella ecaudata*, *Yoldiella sabrina*, and *Yoldiella valettei* (Yoldiidae) from the Southern Ocean. *Polar Biology*, 37(10), 1383-1392. doi:10.1007/s00300-014-1528-4
- Riascos, J. M. (2006). Effects of El Nino-Southern oscillation on the population dynamics of the tropical bivalve *Donax dentifer* from Malaga bay, Colombian Pacific. *Marine Biology*, 148(6), 1283-1293. doi:10.1007/s00227-005-0165-4
- Riascos, J. M., Heilmayer, O., & Laudien, J. (2008). Population dynamics of the tropical bivalve *Cardita affinis* from Málaga Bay, Colombian Pacific related to La Niña 1999–2000. *Helgoland Marine Research*, 62(1), 63-71. doi:10.1007/s10152-007-0083-6
- Riascos, J. M., & Urban, H. J. (2002). Population dynamics of *Donax dentifer* (Vencroida : Donacidae) in Bahia Malaga, Colombian Pacific, during the "El Nino" event 1997/1998. *Revista De Biologia Tropical*, 50(3-4), 1113-1123.
- Richard, G. (1981). *first evaluation of the findings of the growth and production of lagoon and reef molluscs in French Polynesia*. Paper presented at the Proceedings of the Fourth Coral Reef Symposium
- Richardson, C., Seed, R., Al-Roumaihi, E. M. H., & McDonald, L. (1993). Distribution, shell growth and predation of the New Zealand oyster, *Tiostrea (=Ostrea) lutaria* Hutton, in the Menai Strait, North Wales. *Journal of Shellfish Research*, 12(2), 207-214.
- Richardson, C., Seed, R., & Naylor, E. (1990). Use of internal growth bands for measuring individual and population growth rates in *Mytilus edulis* from offshore platforms. *Marine Ecology Progress Series*, 66, 259-265.
- Richardson, C. A., Kennedy, H., Duarte, C. M., Kennedy, D. P., & Proud, S. V. (1999). Age and growth of the fan mussel *Pinna nobilis* from south-east Spanish Mediterranean seagrass (*Posidonia oceanica*) meadows. *Marine Biology*, 133(2), 205-212. doi:10.1007/s002270050459
- Richardson, C. A., Peharda, M., Kennedy, H., Kennedy, P., & Onofri, V. (2004). Age, growth rate and season of recruitment of *Pinna nobilis* (L) in the Croatian Adriatic determined from Mg:Ca and Sr:Ca shell profiles. *Journal of Experimental Marine Biology and Ecology*, 299(1), 1-16. doi:https://doi.org/10.1016/j.jembe.2003.08.012
- Richardson, C. A., Taylor, A. C., & Venn, T. J. (1982). Growth of the Queen Scallop *Chlamys opercularis* in Suspended Cages in the Firth of Clyde. *Journal of the Marine Biological Association of the United Kingdom*, 62(1), 157-169. doi:10.1017/S002531540002018X
- Richardson, M. G. (1979). The ecology and reproduction of the brooding Antarctic bivalve *Lissarca miliaris*. *British Antarctic Survey Bulletin*, 49, 91-115.
- Ridgway, I. D., Richardson, C., Scourse, J., Butler, P., & Reynolds, D. (2012). The population structure and biology of the ocean quahog, *Arctica islandica*, in Belfast Lough, Northern Ireland. *Journal of the Marine Biological Association of the United Kingdom*, 92. doi:10.1017/S0025315411000154
- Ridgway, I. D., Richardson, C. A., Enos, E., Ungvari, Z., Austad, S. N., Philipp, E. E. R., & Csiszar, A. (2011). New Species Longevity Record for the Northern Quahog (=Hard Clam), *Mercenaria mercenaria*. *Journal of Shellfish Research*, 30(1), 35-38. doi:10.2983/035.030.0106



- Robert, G., & Jamieson, G. S. (1986). Commercial fishery data isopleths and their use in offshore sea scallop (*Placopecten magellanicus*) stock evaluations. *Canadian Special Publication of Fisheries and Aquatic Sciences*, 92, 76-82.
- Robinson, R. F., & Richardson, C. A. (1998). The direct and indirect effects of suction dredging on a razor clam (*Ensis arcuatus*) population. *Ices Journal of Marine Science*, 55(5), 970-977. doi:10.1006/jmsc.1998.0356
- Robinson, T. B., Govender, A., Griffiths, C. L., & Branch, G. M. (2007). Experimental harvesting of *Mytilus galloprovincialis*: Can an alien mussel support a small-scale fishery? *Fisheries Research*, 88(1), 33-41. doi:https://doi.org/10.1016/j.fishres.2007.07.005
- Rolfe, M. S. (1973). *Notes on queen scallops and how to catch them*. Ministry of Agriculture Fisheries and Food.
- Saeedi, H., A. Ardalan, A., Kamrani, E., & Kiabi, B. (2010). Reproduction, growth and production of *Amiantis umbonella* (Bivalvia: Veneridae) on northern coast of the Persian Gulf, Bandar Abbas, Iran. *Journal of The Marine Biological Association of The United Kingdom*, 90, 711-718. doi:10.1017/S0025315409991056
- Şahin, C., Düzgüneş, E., Mutlu, C., Aydin, M., & Emiral, H. (1999). Determination of the Growth Parameters of the *Anadara cornea* R. 1844 population by the Bhattacharya Method in the Eastern Black Sea. *Turkish Journal of Zoology*, 23(1), 99 - 106.
- Saloman, C. H., & Taylor, J. L. (1969). Age and growth of large southern quahogs from a Florida estuary. *Proceedings of the National Shellfisheries Association*, 59, 46-51.
- Salzwedel, H. (1979). Salzwedel, H. "Reproduction, growth, mortality, and variations in abundance and biomass of *Tellina fabula* (Bivalvia) in the German Bight in 1975/76. *Veroffentlichungen des Instituts fur Meeresforschung in Bremerhaven*, 18, 111-202.
- Sanchez-Salazar, M. E., Griffiths, C. L., & Seed, R. (1987). The interactive roles of predation and tidal elevation in structuring populations of the edible cockle, *Cerastoderma edule*. *Estuarine, Coastal and Shelf Science*, 25(2), 245-260. doi:https://doi.org/10.1016/0272-7714(87)90125-9
- Sasaki, K. (1982). Growth of the Sakhalin Surf Clam, *Spisula sachalinensis* (SCHRENCK), in Sendai Bay. *Tohoku journal of agricultural research*, 32(4), 168-180.
- Schäffer, F., & Zettler, M. L. (2007). The clam siphon as indicator for growth indices in the soft-shell clam *Mya arenaria*. *Helgoland Marine Research*, 61(1), 9-16. doi:10.1007/s10152-006-0049-0
- Schick, D. F., Shumway, S., & Hunter, M. A. (1988). A comparison of growth rate between shallow water and deep water populations of scallops *Placopecten magellanicus* (Gmelin 1791) in the Gulf of Maine. *American Malacological Bulletin*, 6, 1-8.
- Schmidt, A., Wehrmann, A., & Dittmann, S. (2008). Population dynamics of the invasive Pacific oyster *Crassostrea gigas* during the early stages of an outbreak in the Wadden Sea (Germany). *Helgoland Marine Research*, 62(4), 367. doi:10.1007/s10152-008-0125-8
- Schweers, T., Wolff, M., Koch, V., & Sinsel Duarte, F. (2006). Population dynamics of *Megapitaria squalida* (Bivalvia: Veneridae) at Magdalena Bay, Baja California Sur, Mexico. *Revista De Biologia Tropical*, 54, 1003-1017.
- Seed, R. (1973). Absolute and allometric growth in the mussel, *Mytilus edulis* L. (Mollusca: Bivalvia) *Journal of Molluscan Studies*, 40(5), 343-357. doi:10.1093/oxfordjournals.mollus.a065232
- Sejr, M., & Christensen, P. (2007). Growth, production and carbon demand of macrofauna in Young Sound, with special emphasis on the bivalves *Hiatella arctica* and *Mya truncata*. *Meddelelser Om Grønland*, 58, 121-136.
- Sejr, M. K., Sand, M. K., Jensen, K. T., Petersen, J. K., Christensen, P. B., & Rysgaard, S. (2002). Growth and production of *Hiatella arctica* (Bivalvia) in a high-Arctic fjord (Young Sound, Northeast Greenland). *Marine Ecology Progress Series*, 244, 163-169. doi:10.3354/meps244163



- 1
- 2
- 3 Selin, N., & Selina, M. (1988). Production characteristics of the bivalve mollusc *Callista*
- 4 *brevisiphonata* in Peter the Great Bay, Sea of Japan. *Soviet Journal of Marine Biology*, 14,
- 5 219-223.
- 6
- 7 Selin, N. I. (1980). Size-age structure of settlements of *Crenomytilus grayanus* on differnt grounds in
- 8 Pos'et Bay, Sea of Japan. *Soviet Journal of Marine Biology*, 44-49.
- 9 Selin, N. I. (1988). Size-age structure and growth of the mussel *Mytilus coruscus* in Peter the Great
- 10 Bay, Sea of Japan. *Soviet Journal of Marine Biology*, 14, 284-288.
- 11 Selin, N. I. (1993). Production and Growth of the Bivalve Mollusk *Keenocardium californiense* in the
- 12 Northwestern Part of the Sea of Japan. *Russian Journal of Marine Biology*, 19, 26-33.
- 13 Selin, N. I. (2007). Shell form, growth and life span of *Astarte arctica* and *A. borealis* (Mollusca:
- 14 Bivalvia) from the subtidal zone of northeastern Sakhalin. *Russian Journal of Marine Biology*,
- 15 33(4), 232-237. doi:10.1134/s1063074007040050
- 16
- 17 Selin, N. I. (2008). Distribution, population structure and growth of *Protothaca euglypta* (Sowerby,
- 18 1914) (Bivalvia: Veneridae) from the northwestern part of the East Sea of Russia. *Korean*
- 19 *Journal of Malacology*, 24, 81-87.
- 20 Selin, N. I. (2010). The growth and life span of bivalve mollusks at the northeastern coast of Sakhalin
- 21 Island. *Russian Journal of Marine Biology*, 36(4), 258-269. doi:10.1134/s1063074010040048
- 22 Selin, N. I. (2014). The distribution, size, and age composition, and growth of *Saxidomus purpurata*
- 23 (Sowerby, 1852) (Bivalvia: Veneridae) in Vostok Bay, the Sea of Japan. *Russian Journal of*
- 24 *Marine Biology*, 40(5), 375-382. doi:10.1134/s106307401405006x
- 25
- 26 Selin, N. I., & Latypov, Y. Y. (2011). Size and age composition of *Tridacna crocea* Lamarck, 1819
- 27 (Bivalvia: Tridacnidae) in coastal waters of islands of the Con Dao Archipelago in the South
- 28 China Sea. *Biologiya Morya-Marine Biology*, 37(5), 367-373.
- 29 Selin, N. I., & Lysenko, V. N. (2006). Size and age composition of populations and growth of *Mytilus*
- 30 *trossulus* (Bivalvia: Mytilidae) in the subtidal area of western Kamchatka. *Russian Journal of*
- 31 *Marine Biology*, 32(6), 360. doi:10.1134/s1063074006060058
- 32
- 33 Sephton, T., & Bryan, C. F. (1990). Age and growth rate determinations for the Atlantic surf clam,
- 34 *Spisula solidissima* (Dillwyn, 1817) in Prince Edward Island, Canada. *Journal Shellfish*
- 35 *Research*, 1, 177-185.
- 36 Serchuk, F. M., Wood, P. W., Posgay, J. A., & Brown, B. E. (1979). Assessment and status of sea
- 37 scallop (*Placopecten magellanicus*) populations off the northeast coast of the United States.
- 38 *Proceedings of the National Shellfisheries Association*, 69, 161-191.
- 39 Serchuk, F. M., Wood, P. W., & Rak, R. S. (1982). *Review and assessment of the Georges Bank, Mid-*
- 40 *Atlantic and Gulf of Maine Atlantic sea scallop (Placopecten magellanicus) resources:*
- 41 National Marine Fisheries Service, Northeast Fisheries Center, Woods Hole Laboratory.
- 42 Shafee, M. (1982). Seasonal variation in oxygen consumption rates of the black scallop *Chlamys varia*
- 43 (L.) from Lanveoc (Bay of Brest). *Oceanologica acta Paris*, 5(2), 189-197.
- 44 Shafee, M. (1992). Production estimate of a mussel population *Perna picta* (Born) on the Atlantic
- 45 coast of Morocco. *Journal of Experimental Marine Biology and Ecology*, 163(2), 183-197.
- 46 Shafee, M., & Conan, G. (1984). Energetic parameters of a population of *Chlamys varia* (Bivalvia:
- 47 Pectinidae). *Marine Ecology Progress Series*, 253-262.
- 48 Shafee, M. S. (1979). Ecological energy requirements of the green mussel, *Perna viridis* Linnaeus
- 49 from Ennore estuary, Madras. *Oceanologica Acta*, 2(1), 69-74.
- 50 Shelley, C. C. (1988). Growth rates of *Hippopus hippopus* from Orpheus Island, Great Barrier Reef. In
- 51 J. W. Copland & J. S. Lucas (Eds.), *Giant Clams in Asia and the Pacific* (Vol. 207-212):
- 52 Australian Centre for International Agricultural Research.
- 53 Šiletić, T., & Peharda, M. (2003). Population study of the fan shell *Pinna nobilis* L. in Malo and Veliko
- 54 Jezero of the Mljet National Park (Adriatic Sea). *Scientia Marina*, 67(1), 91-98.
- 55 Silina, A., & Pozdnyakova, L. (1990). Growth of the scallop *Chlamys rosealbus* in the Sea of Japan.
- 56 *Soviet. Journal of Marine Biology*, 16(1), 32-36.
- 57
- 58
- 59
- 60

- Silina, A. V. (2014). Habitat Preferences and growth of *Ruditapes bruguieri* (Bivalvia: Veneridae) at the Northern Boundary of Its Range. *The Scientific World Journal*, 2014, 6. doi:10.1155/2014/235416
- Silva, C. F., Corte, G. N., Yokoyama, L. Q., Abrahao, J. R., & Amaral, A. C. Z. (2015). Growth, mortality, and reproduction of *Tagelus plebeius* (Bivalvia: Solecurtidae) in Southeast Brazil. *Helgoland Marine Research*, 69(1), 1.
- Sims, A. N. (1992). Population dynamics and stock management of the black-lip pearl oyster, *Pinctada margaritifera* (L.), in the Cook Islands, South Pacific. *Marine and Freshwater Research*, 43(6), 1423-1435.
- Singh, Y. T., Krishnamoorthy, M., & Thippeswamy, S. (2011). Population ecology of the wedge clam *Donax faba* (Gmelin) from the Panambur beach, near Mangalore, south west coast of India. *Journal of Theoretical and Experimental Biology*, 7, 171-182.
- Smith, E. B., Scott, K. M., Nix, E. R., Korte, C., & Fisher, C. R. (2000). Growth and condition of seep mussels (*Bathymodiolus childressi*) at a Gulf of Mexico brine pool. *Ecology*, 81(9), 2392-2403. doi:10.1890/0012-9658(2000)081[2392:GACOSM]2.0.CO;2
- Smith, S. D. (2011). Growth and population dynamics of the giant clam *Tridacna maxima* (Röding) at its southern limit of distribution in coastal, subtropical eastern Australia. *Molluscan Research*, 31(1), 37.
- Sola, J. C. (1997). Reproduction, population dynamics, growth and production of *Scrobicularia plana*, da costa (pelecypoda) in the Bidasoa estuary, Spain. *Netherland Journal of Aquatic Ecology*, 30(4), 283-296. doi:10.1007/bf02085872
- Somasekar, M., Sriraman, K., & Kasinathan, R. (1982). Age, Growth and Length-Weight Relationship in the Backwater Oyster *Crassostrea madrasensis* (Preston). *Indian Journal of Marine Sciences*, 11(2), 190-192.
- Soon, T. K., Denil, D. J., & Ransangan, J. (2016). High mortality and poor growth of green mussels, *Perna viridis*, in high chlorophyll-a environment. *Ocean Science Journal*, 51(1), 43-57.
- Steffani, C. N., & Branch, G. M. (2003). Growth rate, condition, and shell shape of *Mytilus galloprovincialis*: responses to wave exposure. *Marine Ecology Progress Series*, 246, 197-209.
- Stern-Pirlot, A., & Wolff, M. (2006). Population dynamics and fisheries potential of *Anadara tuberculosa* (Bivalvia : Arcidae) along the Pacific coast of Costa Rica. *Revista De Biologia Tropical*, 54, 87-100.
- Stockton, W. L. (1984). The biology and ecology of the epifaunal scallop *Adamussium colbecki* on the west side of McMurdo Sound, Antarctica. *Marine Biology*, 78(2), 171-178. doi:10.1007/bf00394697
- Stotz, W. B., & Gonzalez, S. A. (1997). Abundance, growth, and production of the sea scallop *Argopecten purpuratus* (Lamarck 1819): bases for sustainable exploitation of natural scallop beds in north-central Chile. *Fisheries Research*, 32(2), 173-183.
- Strahl, J., Philipp, E., Brey, T., Broeg, K., & Abele, D. (2007). Physiological aging in the Icelandic population of the ocean quahog *Arctica islandica*. *Aquatic Biology*, 1(1), 77-83. doi:10.3354/ab00008
- Sugiura, D., Katayama, S., Sasa, S., & Sasaki, K. (2014). Age and Growth of the Ark Shell *Scapharca Broughtonii* (Bivalvia, Arcidae) in Japanese Waters. *Journal of Shellfish Research*, 33(1), 315-324.
- Sukhotin, A. A., Abele, D., & Portner, H. O. (2006). Ageing and metabolism of *Mytilus edulis*: Populations from various climate regimes. *Journal of Shellfish Research*, 25(3), 893-899.
- Sukhotin, A. A., Strelkov, P. P., Maximovich, N. V., & Hummel, H. (2007). Growth and longevity of *Mytilus edulis* (L.) from northeast Europe. *Marine Biology Research*, 3(3), 155-167. doi:10.1080/17451000701364869
- Swennen, C., Leopold, M. F., & Stock, M. (1985). Notes on growth and behavior of the American razor clam *Ensis directus* in the Wadden Sea and the Predation on it by birds. *Helgolander Meeresuntersuchungen*, 39(3), 255-261.

- 1
- 2
- 3
- 4 Taib, A. M., Madin, J., & Ransangan, J. (2016). Density, recruitment and growth performance of Asian
- 5 green mussel (*Perna viridis*) in Marudu Bay, Northeast Malaysian Borneo, three years after a
- 6 massive mortality event. *Songklanakarin Journal of Science & Technology*, 38(6).
- 7 Takeuchi, S., Suzuki, Y., Takamasu, T., Isomoto, H., & Tamaki, A. (2016). Ecology of the razor clam
- 8 *Solen gordonis* and fishery impact on the population in Sasebo Bay, Kyushu, Japan. *Journal of*
- 9 *Shellfish Research*, 35(4), 785-800.
- 10 Tanabe, K. (1988). Age and growth rate determinations of an intertidal bivalve, *Phacosoma*
- 11 *japonicum*, using internal shell increments. *Lethaia*, 21(3), 231-241.
- 12 Tanabe, K., & Oba, T. (1988). Latitudinal variation in shell growth patterns of *Phacosoma japonicum*
- 13 (Bivalvia: Veneridae) from the Japanese coast. *Marine Ecology Progress Series*, 75-82.
- 14 Taylor, A. C., & Venn, T. J. (2009). Growth of the queen scallop, *Chlamys opercularis*, from the Clyde
- 15 Sea area. *Journal of the Marine Biological Association of the United Kingdom*, 58(3), 687-700.
- 16 doi:10.1017/S0025315400041333
- 17 Tenjing, S., Thippeswamy, S., & Narasimhaiah, N. (2016). A first study on growth, mortality and
- 18 survivorship of the eared horse mussel, *Modiolus auriculatus* (Krauss, 1848) from India.
- 19 Thébault, J., Thouzeau, G., Chauvaud, L., Cantillán, M., & Avendaño, M. (2008). Growth of
- 20 *Argopecten purpuratus* (Mollusca: Bivalvia) on a natural bank in Northern Chile:
- 21 sclerochronological record and environmental controls. *Aquatic Living Resources*, 21(1), 45-
- 22 55.
- 23 Theisen, B. F. (1973). The growth of *Mytilus edulis* L. (Bivalvia) from Disko and Thule District,
- 24 Greenland. *Ophelia*, 12(1&2), 59-77.
- 25 Theisen, B. F. (1975). Growth parameters of *Mytilus edulis* L. (Bivalvia) estimated from tagging data.
- 26 *Meddelelser fra Danmarks Fiskeri og Havundersoegelser*.
- 27 Thippeswamy, S., & Joseph, M. M. (1991). Population Selection-Strategies in the Wedge Clam,
- 28 *Donax-Incaratus* (Gmelin) from Panambur Beach, Mangalore. *Indian Journal of Marine*
- 29 *Sciences*, 20(2), 147-151.
- 30 Thomas, S., & Nasser, M. (2009). Growth and population dynamics of short-neck clam *Paphia*
- 31 *malabarica* from Dharmadam estuary, North Kerala, southwest coast of India. *Journal of the*
- 32 *Marine Biological Association of India*, 51(1), 87-92.
- 33 Thompson, R. J. (1984). Production, reproductive effort, reproductive value and reproductive cost in
- 34 a population of the blue mussel *Mytilus edulis* from a subarctic environment. *Marine Ecology*
- 35 *Progress Series*, 16, 249-257.
- 36 Thorarinsdóttir, G. G., & Jacobson, L. D. (2005). Fishery biology and biological reference points for
- 37 management of ocean quahogs (*Arctica islandica*) off Iceland. *Fisheries Research*, 75(1), 97-
- 38 106. doi:https://doi.org/10.1016/j.fishres.2005.04.010
- 39 Thouzeau, G., Robert, G., & Smith, S. J. (1991). Spatial Variability in Distribution and Growth of
- 40 Juvenile and Adult Sea Scallops *Placopecten magellanicus* (Gmelin) on Eastern Georges Bank
- 41 (Northwest Atlantic). *Marine Ecology Progress Series*, 74(2-3), 205-218.
- 42 Tunberg, B. (1983a). Growth of *Dosinia exoleta* (L) (Bivalvia) in Raunefjorden, Western Norway.
- 43 *Sarsia*, 68(1), 41-45.
- 44 Tunberg, B. (1983b). Population-Structure, Size Distribution, and Shell Growth of *Dosinia lupinus* (L)
- 45 (Bivalvia) in Raunefjorden, Western Norway, with Biometrical Comparison to *Dosinia-Exoleta*
- 46 (L). *Sarsia*, 68(1), 33-40.
- 47 Turra, A., Petracco, M., Amaral, A. C. Z., & Denadai, M. R. (2014). Temporal variation in life-history
- 48 traits of the clam *Tivela mactroides* (Bivalvia: Veneridae): Density-dependent processes in
- 49 sandy beaches. *Estuarine Coastal and Shelf Science*, 150, 157-164.
- 50 Turra, A., Petracco, M., Amaral, A. C. Z., & Denadai, M. R. (2015). Population biology and secondary
- 51 production of the harvested clam *Tivela mactroides* (Born, 1778) (Bivalvia, Veneridae) in
- 52 Southeastern Brazil. *Marine Ecology*, 36(2), 221-234. doi:10.1111/maec.12137
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- Urban, H.-J. (2000). Culture potential of the pearl oyster (*Pinctada imbricata*) from the Caribbean.: I. Gametogenic activity, growth, mortality and production of a natural population. *Aquaculture*, 189(3-4), 361-373.
- Urban, H. J. (1996). Population dynamics of the bivalves *Venus antiqua*, *Tagelus dombeii*, and *Ensis macha* from Chile at 36 degrees South. *Journal of Shellfish Research*, 15(3), 719-727.
- Urban, H. J. (1998). Description and management of a clam fishery (*Gari solida*, Psammobiidae) from Bahia Independencia, Peru (14 degrees S). *Fisheries Research*, 35(3), 199-207. doi:10.1016/s0165-7836(98)00075-7
- Urban, H. J., & Campos, B. (1994). Population-Dynamics of the Bivalves *Gari solida*, *Semele solida* and *Protothaca thaca* from a Small Bay in Chile at 36-Degrees-S. *Marine Ecology-Progress Series*, 115(1-2), 93-102.
- Urban, H. J., & Mercuri, G. (1998). Population dynamics of the bivalve *Laternula elliptica* from Potter cove, King George Island, South Shetland islands. *Antarctic Science*, 10(2), 153-160.
- Urban, H. J., & Tarazona, J. (1996). Effects of El Nino Southern Oscillation on the population dynamics of a *Gari solida* population (Bivalvia:Psammobiidae) from Bahia Independencia, Peru. *Marine Biology*, 125(4), 725-734. doi:10.1007/bf00349255
- Urban, H. J., & Tesch, C. (1996). Aspects of the population dynamics of six bivalve species from Southern Chile. Results of the 'Victor Hensen' cruise to the Magellan Strait and the Beagle Channel in October/November 1994. *Archive of Fishery and Marine Research*, 44(2), 243-256.
- Urrutia, M. B., Ibarrola, I., Iglesias, J. I. P., & Navarro, E. (1999). Energetics of growth and reproduction in a high-tidal population of the clam *Ruditapes decussatus* from Urdaibai Estuary (Basque Country, N. Spain). *Journal of Sea Research*, 42(1), 35-48. doi:10.1016/s1385-1101(99)00017-9
- Ursin, E. (1956). Distribution and growth of the queen scallop *Chlamys opercularis* (Lamellibranchiata) in Danish and Faroese waters. *Meddelelser fra Danmarks Fiskeri og Havundersoegelser*, 1, 1 - 31.
- Vahl, O. (1981). Energy transformations by the Iceland scallop, *Chlamys islandica* (OF Müller), from 70 NI The age-specific energy budget and net growth efficiency. *Journal of Experimental Marine Biology and Ecology*, 53(2-3), 281-296.
- Vakily, J. (1992). *Determination and comparison of bivalve growth, with emphasis on Thailand and other tropical areas* (Vol. 801): WorldFish.
- Valero, J. (1999). Variación estacional, espacial e interannual en el crecimiento de vieira patagónica (*Zygochlamys patagonica*) en la plataforma argentina. *Informe Final-Contrato IICA*.
- Velarde, A. A., Flye-Sainte-Marie, J., Mendo, J., & Jean, F. (2015). Sclerochronological records and daily microgrowth of the Peruvian scallop (*Argopecten purpuratus*, Lamarck, 1819) related to environmental conditions in Paracas Bay, Pisco, Peru. *Journal of Sea Research*, 99, 1-8.
- Velazquez-Abunader, I., Lopez-Rocha, J. A., Arellano-Martinez, M., Ceballos-Vazquez, B. P., & Cabrera, M. A. (2016). Estimation of growth parameters in a wild population of lion-paw scallop (*Nodipecten subnodosus*) in Bahia de Los Angeles, Baja California, Mexico. *Hidrobiologica*, 26(1), 133-142.
- Vélez, A., Venables, B. J., & Fitzpatrik, L. (1985). Growth production of the tropical beach clam *Donax denticulatus* (Tellinidae) in eastern Venezuela. *Caribbean Journal of Science*, 21, 63-73.
- Veloso, V., Moreira, J., & Troncoso, J. S. (2007). Annual dynamics of bivalve populations in muddy bottoms of the Ensenada de Baiona (Galicia, NW Iberian Peninsula). *Iberus*, 25, 1-10.
- Ventilla, R. F. (1982). The Scallop Industry in Japan. In J. H. S. Blaxter, F. S. Russell, & M. Yonge (Eds.), *Advances in Marine Biology* (Vol. 20, pp. 309-382): Academic Press.
- Verdelhos, T., Neto, J. M., Marques, J. C., & Pardal, M. A. (2005). The effect of eutrophication abatement on the bivalve *Scrobicularia plana*. *Estuarine Coastal and Shelf Science*, 63(1-2), 261-268.



- 1
- 2
- 3 Verginelli, R., & Prieto, A. (1991). Produccion secundaria de *Pinctada imbricata* (Roding, 1798)
- 4 (Pterioda:Pteriidae) en una poblacion del Golfo de Cariaco, Venezuela. *Acta Cient. Venez.*,
- 5 42, 138-144.
- 6
- 7 Verween, A., Vincx, M., & Degraer, S. (2006). Growth patterns of *Mytilopsis leucophaeata*, an
- 8 invasive biofouling bivalve in Europe. *Biofouling*, 22(4), 221-231.
- 9 doi:10.1080/08927010600816401
- 10 Vincent, B., Brassard, C., & Harvey, M. (1987). Variations de la croissance de la coquille, et de la
- 11 structure d'âge du bivalve *Macoma balthica* (L.) dans une population intertidale de l'estuaire
- 12 du Saint-Laurent (Québec). *Canadian Journal of Zoology*, 65(8), 1906-1916. doi:10.1139/z87-
- 13 291
- 14 Vuorinen, I., Antsulevich, A. E., & Maximovich, N. V. (2002). Spatial distribution and growth of the
- 15 common mussel *Mytilus edulis* L. in the archipelago of SW-Finland, northern Baltic Sea.
- 16 *Boreal environment research*, 7(1), 41-52.
- 17
- 18 Walker, R. L., & Heffernan, P. B. (1994). Age, growth rate, and size of the southern surfclam, *Spisula*
- 19 *solidissima similis* (Say, 1822). *Journal of Shellfish Research*, 13(2), 433-441.
- 20 Walker, R. L., & Tenore, K. R. (1984). The distribution and production of the hard clam, *Mercenaria*
- 21 *mercenaria*, in Wassaw Sound, Georgia. *Estuaries*, 7(1), 19.
- 22
- 23 Waloszek, D. (1991). *Chlamys patagonica* (King & Broderip, 1832), a long "neglected" species from
- 24 the shelf off the Patagonian Coast. *An international compendium of scallop biology and*
- 25 *culture*, 256-263.
- 26
- 27 Warwick, R. M., George, C. L., & Davies, J. R. (1978). Annual Macrofauna Production in a *Venus*
- 28 Community. *Estuarine and Coastal Marine Science*, 7(3), 215-241.
- 29 Weinberg, J. R., & Helser, T. E. (1996). Growth of the Atlantic surfclam, *Spisula solidissima*, from
- 30 Georges Bank to the Delmarva Peninsula, USA. *Marine Biology*, 126(4), 663-674.
- 31 doi:10.1007/bf00351333
- 32
- 33 Wendell, F., Demartini, J. D., Dinnel, P., & Siecke, J. (1976). Ecology of Gaper or Horse Clam, *Tresus-*
- 34 *capax* (Gould 1850) (Bivalvia-Mactridae), in Humboldt Bay, California. *California Fish and*
- 35 *Game*, 62(1), 41-64.
- 36
- 37 Weymouth, F. W., McMillin, H. C., United, S., & Bureau of, F. (1931). *Relative growth and mortality of*
- 38 *the Pacific razor clam (Siliqua patula, Dixon) and their bearing on the commercial fishery.*
- 39 Washington, D.C.: U.S. G.P.O.
- 40
- 41 Weymouth, F. W., & Thompson, S. H. (1930). Thompson, S.H. 1930 The age and growth of the Pacific
- 42 cockle (*Cardium corbis*, Martyn). *Bulletin of the Bureau of Fisheries*, 46, 633-641.
- 43
- 44 Williams, M., & Dredge, M. (1981). Growth of the saucer scallop, *Amusium japonicum* balloti Habe,
- 45 in central eastern Queensland. *Marine and Freshwater Research*, 32(4), 657-666.
- 46
- 47 Wilson, J. G. (1997). Long-term changes in density, population structure and growth rate of *Tellina*
- 48 *tenuis* from Dublin Bay, Ireland. *Oceanologica Acta*, 20(1), 267-274.
- 49
- 50 Winther, U., & Gray, J. S. (1985). The Biology of *Mya arenaria* (Bivalvia) in the Eutrophic Inner
- 51 Oslofjord. *Sarsia*, 70(1), 1-9.
- 52
- 53 Wolf, B., & White, R. (1995). Age and growth of the queen scallop, *Equichlamys bifrons*, in the
- 54 D'Entrecasteaux Channel and Huon River Estuary, Tasmania. *Marine and Freshwater*
- 55 *Research*, 46(8), 1127-1135.
- 56
- 57 Wolfe, D. A., & Petteway, E. N. (1968). Growth of *Rangia cuneata* Gray. *Chesapeake Science*, 9(2), 99-
- 58 102. doi:10.2307/1351251
- 59
- 60 Wolff, M. (1985). Abundancia masiva y crecimiento de pre-adultos de la concha de abanico peruana
- (*Argopecten purpuratus*) en la zona de Pisco bajo condiciones de El Niño 1983.
- Xavier, B. M., Branch, G. M., & Wieters, E. (2007). Abundance, growth and recruitment of *Mytilus*
- galloprovincialis* on the west coast of South Africa in relation to upwelling. *Marine Ecology*
- Progress Series*, 346, 189-201.

Yamashiro, C., & Mendo, J. (1988). Crecimiento de la concha de abanico (*Argopecten purpuratus*) en la Bahía Independencia, Pisco, Perú. *Revista de Biología Marina y Oceanografía*, 42(3) 275-285. Doi: 10.4067/S0718-19572007000300008

Yap, W. G. (1977). Population biology of the Japanese little-neck clam, *Tapes philippinarum*, in Kaneohe Bay, Oahu, Hawaiian Islands. *Pacific Science*, 31(3), 223-244.

Yoo, S. K., Ryu, H.-Y., & PARK, K.-Y. (1981). The growth of the cultured scallop, *Patinopecten yessoensis*. *Korean Journal of Fisheries and Aquatic Sciences*, 14(4), 221-226.

Yoon, H.-S., An, Y.-K., Kim, S.-T., & Choi, S.-D. (2011). Age and growth of the short necked *Ruditapes philippinarum* on the south coast of Korea. *The Korean Journal of Malacology*, 27(1), 1-7.

Zaidman, P. C., & Morsan, E. (2015). Growth variability in a metapopulation: The case of the southern geoduck (*Panopea abbreviata*). *Fisheries Research*, 172, 423-431. doi:10.1016/j.fishres.2015.08.011


Zeichen, M. M., Agnesi, S., Mariani, A., Maccaroni, A., & Ardizzone, G. D. (2002). Biology and population dynamics of *Donax trunculus* L. (Bivalvia : Donacidae) in the South Adriatic Coast (Italy). *Estuarine Coastal and Shelf Science*, 54(6), 971-982.

Zeinalipour, M., Kiabi, B. H., Shokri, M. R., & Ardalan, A. A. (2014). Population dynamic and distribution of *Barbatia decussata* (Bivalvia: Arcidae) on rocky intertidal shores in the northern Persian Gulf (Iran). *Tropical Zoology*, 27(3), 73-87. doi:10.1080/03946975.2014.944381

Zotin, A. A., & Ozernyuk, N. D. (2004). Growth Characteristics of the Common Mussel *Mytilus edulis* from the White Sea. *Biology Bulletin of the Russian Academy of Sciences*, 31(4), 377-381. doi:10.1023/B:BIBU.0000036942.56020.d2


Supplementary Appendix 2

Figure S2.1




**Figure S2.1.** Overall growth performance in each biogeographical Province. In each case, the median is indicated at the midpoint, the upper and lower quartiles are indicated by the hinges, lines represent the spread and open circles indicate outliers. Data points are superimposed.

Figure S2.2




**Figure S2.2** Relationship of temperature (derived from World Ocean Atlas 2013) and latitude based on each data point used in this study. Geographic realms are indicated by colour (see inset legend).




**Figure S2.3**

**Figure S2.3** Relationship of particulate organic carbon and latitude for each data point used in this study. Geographic realms are indicated by colour (see inset legend).




**Figure S2.4** Relationship of overall growth performance in the subclasses Pteriomorpha (blue), Heterodonta (red) and Protobranchia (green) with a) temperature; b) latitude; c) Particulate Organic Carbon.

Figure S2.5




**Figure S2.5** Relationship of overall growth performance at the taxonomic resolution of family (indicated below panels) within the subclasses Pteriimorphia (top panel), and Heterodonta (bottom panel) with Temperature, Latitude, and Particulate Organic Carbon.

Figure S2.6



**Figure S2.6.** Linear regressions of inverse temperature (Kelvins) and natural log of Overall Growth Performance, representing the Arrhenius Relationship in four biogeographic latitude zones a) Tropical; b) Temperate; c) Arctic; d) Antarctic.



**Figure S2.7**

**Figure S2.7.** Linear regressions of inverse temperature (Kelvins) and natural log of Overall Growth Performance, representing the Arrhenius Relationship, in twelve biogeographic realms.